


# Regulation of flowering time by FVE, a retinoblastoma-associated protein

Israel Ausín<sup>1,3</sup>, Carlos Alonso-Blanco<sup>1-3</sup>, José A Jarillo<sup>2</sup>, Leonor Ruiz-García<sup>1</sup> & José M Martínez-Zapater<sup>1,2</sup>

The initiation of flowering in plants is controlled by environmental and endogenous signals<sup>1,2</sup>. Molecular analysis of this process in *Arabidopsis thaliana* indicates that environmental control is exerted through the photoperiod and vernalization pathways, whereas endogenous signals regulate the autonomous and gibberellin pathways. The vernalization and autonomous pathways converge on the negative regulation of *FLC*<sup>3,4</sup>, a gene encoding a MADS-box protein that inhibits flowering<sup>3,4</sup>. We cloned *FVE*, a component of the autonomous pathway that encodes AtMSI4, a putative retinoblastoma-associated protein. *FVE* interacted with retinoblastoma protein

in immunoprecipitation assays, and *FLC* chromatin was enriched in acetylated histones in *fve* mutants. We conclude that *FVE* participates in a protein complex repressing *FLC* transcription through a histone deacetylation mechanism. Our data provide genetic evidence of a new developmental function of these conserved proteins and identify a new genetic mechanism in the regulation of flowering.

Many plants require prolonged exposure to near-freezing temperatures (vernalization) to flower. This reversible delay is associated in *A. thaliana* with the expression of the *FLC* gene<sup>3,4</sup>. In early flowering



**Figure 1** Phenotype and genotype of *fve* mutants. (a) Col and *fve-3* plants grown for 4 weeks under long-day (16-h) or 6 weeks under short-day (8-h) photoperiods with (V+) or without (V-) 8 weeks of vernalization at 4 °C. (b) Total number of leaves produced by wild-type and *fve* mutant plants under different environmental conditions. Plants were grown under long days or short days without vernalization (white columns) or after vernalization for 4 weeks (light gray columns) or 12 weeks (dark gray columns). (c) Effect of *fve* mutations on *FVE* and *FLC* expression in 12-d-old vegetative plants. Hybridization with an 18S ribosomal probe is shown as loading control. The rubisco large subunit (*rbcL*) stained with Ponceau is shown as protein loading control. (d) Structure of *FVE* gene (At2g19520) and protein. Exons are shown as boxes and introns as lines. The predicted translation start (ATG) and stop (TAA) codons are indicated. The 5' and 3' end untranslated regions are boxed in black. Mutations in the *fve* alleles are shown in the upper part, capital letters correspond to the wild-type or mutated nucleotides. In the protein schematic, gray boxes represent WD repeats, the narrow black box represents a putative nuclear localization signal and the asterisk indicates a putative retinoblastoma-binding motif.

<sup>1</sup>Departamento de Genética Molecular de Plantas, Centro Nacional de Biotecnología, Consejo Superior de Investigaciones Científicas, Campus de la Universidad Autónoma de Madrid, Cantoblanco, 28049 Madrid, Spain. <sup>2</sup>Departamento de Biotecnología, Instituto Nacional de Investigación y Tecnología Agraria y Alimentaria, Carretera de A Coruña, km 7, 28040, Madrid, Spain. <sup>3</sup>These authors contributed equally to this work. Correspondence should be addressed to J.M.M.-Z. (zapater@cnb.uam.es).

plants that do not require vernalization, *FLC* is expressed at low levels because these plants lack a functional allele at the *FRI* locus that encodes an *FLC* activator<sup>5</sup>. Genes of the flowering autonomous pathway, such as *FCA*, *FPA*, *FVE*, *FY*, *LD* and *FLD* also restrict *FLC* expression in early-flowering plants<sup>6,7</sup>. Mutations in these genes increase levels of *FLC* mRNA, resulting in a photoperiod-independent flowering delay. The increased *FLC* expression and late flowering of plants with mutations of the autonomous pathway can be reverted by vernalization<sup>3,4</sup>, and *flc* null mutations completely correct the late flowering of plants with mutations of this pathway<sup>8</sup>. These effects indicate that the primary function of the autonomous pathway is the negative regulation of *FLC*. Thus, *FLC* integrates flowering signals from the vernalization and autonomous pathways. Molecular identification of several genes in the autonomous pathway suggested potential roles as regulators of *FLC* expression based on the function assigned to the deduced proteins<sup>9–12</sup>. But this information has not provided an understanding of the molecular mechanisms involved in the regulation of *FLC* expression.


*FVE* is a classical flowering time locus of the autonomous pathway<sup>6,13,14</sup>, as *fve* mutants flowered later than wild-type plants in any photoperiod condition (Fig. 1). The delay in flowering time caused by the *fve* mutations was completely corrected by exposure to 4 °C for 12 weeks (Fig. 1b). In addition, *fve* mutations caused an increase in the steady state level of the *FLC* transcript (Fig. 1c), which can be reverted by vernalization<sup>3,4</sup>. For further insight into the regulation of flowering time by the autonomous pathway, we cloned the gene


*FVE* with a map-based approach (Supplementary Fig. 1 online). A genomic sequence on chromosome 2 previously annotated as *AtMSI4* (ref. 15) was identified as *FVE* because it carried mutations in four *fve* alleles (Fig. 1d). In addition, *fve-1* and *fve-3* plants were complemented with a *FVE* genomic fragment (Fig. 2a). *FVE* encodes a predicted protein of 507 amino acids with six WD repeat domains that are frequently found in eukaryotic proteins involved in basic cell regulatory processes<sup>16</sup>. The predicted protein also contains one stretch of basic residues (amino acids 20–30) that resembles a nuclear localization signal<sup>17</sup> (Fig. 1d). Several independent transgenic lines homozygous with respect to a construct carrying a translational fusion of *FVE* to green fluorescent protein (GFP) under the control of *FVE* regulatory sequences showed nuclear GFP signal in most seedling cells (Fig. 2b).

We analyzed the spatial and temporal expression of *FVE* at the transcript and protein level (Fig. 2). *FVE* RNA and protein were detected in all analyzed vegetative organs and in reproductive organs at higher levels (Fig. 2c). Transcript and protein levels were correlated, indicating that the amount of *FVE* protein is mainly determined by transcriptional regulation. We also analyzed *FVE* expression in response to different photoperiod or vernalization treatments (Fig. 2d) and in *FRI-Sf2*, *ld*, *fpa*, *fca* and *fy* plants with mutations of the autonomous pathway (Fig. 2e) but found no significant differences. Therefore, *FVE* expression is not strongly regulated during plant development, nor is it affected by mutations in other genes of the pathway or by photoperiod or low temperatures.

**Figure 2** Molecular characterization of *FVE*.

(a) Complementation of *fve* mutants. Flowering time under long-day photoperiod of two representative *fve-1* and *fve-3* transgenic lines, homozygous with respect to a *Ler* 5.5-kb *FVE* genomic construct. (b) Nuclear localization of *FVE*-GFP fusion protein. DAPI was used for nuclear staining. (c) *FVE* expression in different organs of *Col* plants. R, roots; RL, rosette leaves; CL, cauline leaves; St, main stems; Fl, flower buds; Fr, developing fruits. (d) *FVE* transcript expression in *Ler* and *Col* grown in different environmental conditions. LD, long day; SD, Short day; V+, plants vernalized for 4 weeks; V-, unvernallized plants. (e) *FVE* transcript expression in flowering mutants of the autonomous pathway. (f) Flowering phenotype and *FVE* expression under long-day photoperiod in transgenic plants overexpressing *FVE*. 350X1, 350X2 and 350X3 are three representative *Ler* transgenic lines and 350X4, 350X5 and 350X6 are three representative *Col* transgenic lines, carrying a single homozygous insertion of *FVE* cDNA under the 35S promoter. 350X1 and 350X4 are cosuppressed lines. DOX1, DOX2 and DOX3 are three representative *Ler* transgenic lines and DOX4, DOX5 and DOX6 are three representative *Col* transgenic lines homozygous with respect to a single insertion of a 6.5-kb *FVE* genomic construct. In a and f, light gray bars show *Ler* genetic background lines and dark gray bars show *Col* background lines. In d–f, RNA and proteins were extracted from 14-d-old rosettes. Hybridization with an 18S ribosomal probe and the rubisco large subunit (rbCL) stained with Ponceau are shown as RNA and protein loading controls, respectively.





**Figure 3** Interaction between FVE and the maize retinoblastoma protein ZmRBR1. (a) An immunoprecipitation assay (IP) using protein extracts prepared from seedlings of null mutant *fve-3* and of the 35S:*FVE* line 35OX5, precipitated with antibody to FVE and hybridized in western blots (WB) with antibody to FVE. (b) An immunoprecipitation (IP) assay in which recombinant ZmRBR1 protein was added to the 35S:*FVE* and *fve-3* protein crude extracts and incubated with antibody to FVE. The proteins precipitated in this way were hybridized in western blots (WB) with antibody to ZmRBR1.


Two strategies of genetic manipulation of *FVE* expression confirmed the correlation between protein amount and transcript levels (Fig. 2f). First, homozygous transgenic plants overexpressing the *FVE* cDNA from a 35S promoter showed several times more *FVE* transcript and protein. A high proportion of those transgenic plants (7 of 15 lines) showed cosuppression of *FVE* expression and had phenotypes similar to *fve* mutants (Fig. 2f). Second, increasing the number of genomic *FVE* copies in Landsberg *erecta* (*Ler*) or Columbia (*Col*) transgenic plants caused comparable increases in transcript and protein levels, but no cosuppression was observed. These plants had either a wild-type phenotype or a small reduction in their flowering time when grown under either long days (Fig. 2f) or short days (data not shown). This indicates that *FVE* overexpression is not sufficient to significantly alter flowering time, even under limiting conditions of flowering induction. Therefore, FVE concentration in the cell seems not to be a limiting factor and FVE function might require other protein factors.

*FVE* (or *AtMSI4*) is a plant homolog of yeast MSI (multicopy suppressor of *IRA1*) and mammalian retinoblastoma-associated proteins RbAp46 and RbAp48 (ref. 15). The *A. thaliana* genome contains a small gene family of five MSI-like genes<sup>18–20</sup>. Database searches showed that MSI genes are present in all sequenced eukaryotic genomes and overall, plant genomes have more MSI genes than genomes from other kingdoms. A comparative phylogenetic analysis of *A. thaliana* MSI-like proteins with other similar proteins from plants, yeasts and animals arranged *A. thaliana* MSI proteins in two

main clades<sup>20</sup> (Supplementary Fig. 2 online). *AtMSI1*, *AtMSI2* and *AtMSI3* are more closely related to animal sequences than the other *A. thaliana* proteins and, together with other plant proteins, cluster in a large heterogeneous clade. *FVE* (*AtMSI4*) and *AtMSI5* share a high similarity (75% identity) and are grouped in a clade that includes only plant MSI-like proteins. All these proteins are more similar to one another than to other MSI-like proteins of the same plant species. Hence, divergence among plant MSI-like proteins predates the monocot-dicot divergence. Plant-specific proteins might have been recruited along plant evolution to carry out functions more specifically related to plant biology. Moreover, the specific and distinct phenotypes of *AtMSI1* antisense transgenic plants<sup>20,21</sup> and *fve* (*AtMSI4*) null mutants suggest that they are not functionally redundant, nor are they redundant with other *A. thaliana* MSI-like proteins.

MSI or retinoblastoma-associated proteins are present in protein complexes involved in chromatin assembly and histone modification<sup>20,21</sup>. In *A. thaliana*, *AtMSI1* is associated with the CAF complex<sup>22</sup> and with PcG complexes, such as MEDEA<sup>21</sup>. Additionally, MSI-like proteins associate with retinoblastoma in animal and plant systems, and ZmRbAp1, the maize *FVE* ortholog, interacts *in vitro* with maize retinoblastoma protein ZmRBR1 (ref. 23). As *FVE* possesses one putative retinoblastoma-binding motif (L-X-C-X-D)<sup>24</sup> in the fourth WD repeat (Fig. 1c), we analyzed the interaction between *FVE* and retinoblastoma. Yeast two-hybrid assays did not show significant interactions, probably owing to intrinsic limitations of the system when dealing with transcriptional repressors. For this reason, we carried out immunoprecipitation assays to determine whether *FVE* associates with the ZmRBR1. We added recombinant ZmRBR1 to protein crude extracts from *Col* wild-type plants, null *fve-3* mutants and transgenic *A. thaliana* plants overexpressing *FVE*. ZmRBR1 was immunoprecipitated from extracts of plants overexpressing *FVE* but not of *fve-3* plants, indicating an interaction between *FVE* and ZmRBR1 (Fig. 3) and suggesting that *FVE* could be part of retinoblastoma-containing complexes in *A. thaliana*. Furthermore, ZmRBR1 was not immunoprecipitated from extracts of wild-type *Col* plants (data not shown), probably because all the *FVE* protein in wild-type plants is part of stable protein complexes and not accessible to ZmRBR1.

In yeast and animal systems, retinoblastoma proteins can function as transcriptional repressors by recruiting HD1/RPD3 histone deacetylases (HDACs). Retinoblastoma-associated proteins similar to *FVE* are components of such complexes<sup>25</sup>. Because *FVE* function is required for *FLC* repression, we speculated that *FVE* could repress *FLC* expression by participating in HDAC complexes modifying the chromatin structure of *FLC*. To investigate this hypothesis, we analyzed histone acetylation in *FLC* using chromatin immunoprecipitation (ChIP) assays (Fig. 4). Chromatin of *fve-1*,



**Figure 4** Effect of *fve* mutations on *FLC* histone acetylation by ChIP analysis. (a) *FLC* genomic region analyzed using antibodies to acetylated H3 and H4. The gray box corresponds to the promoter region, white boxes to exons and black box to the first intron. The different *FLC* fragments analyzed by semi-quantitative PCR are represented and numbered below. (b) PCR products after 28 cycles of *Ler* wild type and *fve-1* mutant using DNA purified from chromatin immunoprecipitated with antibodies against acetylated H3 (ACh3). *ACTIN2* was amplified as control for DNA quantification.

*fve-3*, *fca-1*, *Ler* and *Col* plants was immunoprecipitated using antibodies against acetylated histones H3 or H4. We amplified ten DNA fragments spanning the promoter, first exon and first intron of *FLC* (Fig. 4a) from the precipitated chromatin. *FLC* sequences were consistently more abundant in precipitated chromatin from *fve* mutants than from wild-type plants in three independent experiments (Fig. 4b; DNA from *fve-1* and *fve-3* mutants was  $1.9 \pm 0.2$  and  $1.9 \pm 0.3$  times more abundant than that from respective wild-type plants). Comparable but smaller enrichment was observed for acetylated histone H4. In contrast, the same sequences had similar abundances in plants with another mutation of the autonomous pathway, *fca-1*, and in *Ler* plants (mean relative *fca-1/Ler* value =  $1.17 \pm 0.07$ ). Thus, in *fve* mutants, *FLC* chromatin is enriched in acetylated histones, indicating that *FVE* affects histone acetylation at *FLC*. We conclude that *FVE* is required for histone deacetylation of the *FLC* chromatin in *A. thaliana*.

MSI proteins are biochemically well characterized in several organisms, where they have been involved in the negative regulation of cell cycle and cell differentiation<sup>21,25</sup>, but their developmental functions are poorly understood. The identification of *FVE* as a member of a plant-specific clade of retinoblastoma-associated proteins provides the first genetic evidence to our knowledge of a developmental function for this clade of MSI-like proteins. Furthermore, the participation of *FVE* in the negative regulation of *FLC* transcription through histone deacetylation characterizes a new genetic mechanism in the regulation of flowering.

While this manuscript was under review, the gene *FLD* of the autonomous pathway was shown to encode a homolog of a protein found in mammal HDAC complexes<sup>26</sup>. Similar to *fve*, *fld* mutants show hyperacetylation of histones in *FLC* chromatin. These findings further support the involvement of histone acetylation in the regulation of flowering time in *A. thaliana*.

## METHODS

**Plant materials, growth conditions and measurement of flowering time.** We used the following *A. thaliana* strains: the wild-type laboratory accessions *Ler* and *Col*; the *fve-1* and *fve-2* mutated alleles in *Ler* background, generated by ethyl methane sulfonate mutagenesis<sup>6</sup>; the fast neutron-induced *fve-3* and *fve-4* mutated alleles in *Col* background; the *fca-1*, *fpa-1* and *fy-1* flowering mutations of the autonomous pathway in a *Ler* genetic background<sup>6</sup>; and the *ld-1* mutation and the *FRI-Sf2* allele from Saint Feliu accession introgressed in *Ler* background<sup>27</sup>. We measured flowering time as the total number of leaves (rosette and cauline leaves) developed by a plant<sup>6</sup>.

**Molecular characterization of *FVE*.** We obtained genomic constructs, over-expression cDNA constructs and *FVE*-GFP fusion constructs using standard procedures as detailed in **Supplementary Methods** online. We transformed *A. thaliana* plants by the floral dip method. We analyzed *FVE* mRNA expression by hybridization of northern blots containing 30  $\mu$ g of total RNA isolated by standard procedures with the full length *FVE* cDNA. Similarly, we analyzed *FLC* expression using as probe an *EcoRI-SphI* 700-bp fragment of the *FLC* cDNA<sup>3</sup>. Northern blots were hybridized with an 18S ribosomal probe as loading control.

We raised rabbit polyclonal antibodies against a chemically synthesized *FVE* peptide corresponding to a sequence of 26 N-terminal amino acids present only in *FVE* but not in other *A. thaliana* MSI proteins. This antibody recognized a protein of ~56 kDa at equivalent levels in wild-type and *fve-1* plants but not in *fve-3* plants, showing its *FVE* specificity. We used this antibody to analyze *FVE* protein expression by hybridizing western blots containing ~40  $\mu$ g of total proteins extracted by standard procedures, resolved on 8% acrylamide SDS gels and transferred to Biotrace NT nitrocellulose membranes (Pall, Gelman). We detected proteins with SuperSignal West Pico Chemiluminescence Substrate kit (Pierce).

**Interaction between *FVE* and retinoblastoma protein.** We carried out immunoprecipitation assays as described<sup>28</sup> using *A. thaliana* protein extracts prepared from 12-d-old to 14-d-old seedlings grown under long-day photoperiod conditions in solid nutrient medium. We prepared extracts by homogenizing seedlings in ice-cold buffer containing 50 mM Tris Cl, 150 mM NaCl, 0.5% Nonidet P-40, 1 mM phenylmethylsulfonyl fluoride and 1 $\times$  protease inhibitor cocktail. We precleared extracts by spinning for 15 min in a microcentrifuge. Before immunoprecipitation, we incubated 1 mg of extract with 50  $\mu$ l of protein A-agarose (SIGMA) for 2 h at 4  $^{\circ}$ C with gentle mixing followed by brief centrifugation. We then added a total of 10  $\mu$ l of antibody to *FVE* to the precleared extract and incubated it for 12 h at 4  $^{\circ}$ C. To study *in vitro* the putative interaction between recombinant ZmRBR1 and *FVE*, we added recombinant ZmRBR1 (ref. 29) to precleared protein extract isolated from the transgenic line 35OX5 (overexpressing *FVE* under the 35S promoter) and *fve-3* plants previously incubated with antibody to *FVE*. We collected protein complexes between ZmRBR1 and AtFVE after adding 50  $\mu$ l of protein A-agarose and washed them three times for 15 min in 1 ml of ice-cold washing buffer. We analyzed the immunoprecipitated proteins by western blotting using the antibody to *FVE* or a rabbit antiserum against the maize retinoblastoma protein. The latter was derived against the recombinant ZmRBR1 A/B pocket<sup>25</sup>.

**ChIP assays and PCR.** We carried out ChIP assays as described<sup>30</sup> with minor modifications. We collected 10-d-old *Ler*, *fve-1*, *fca-1*, *Col* and *fve-3* plants, at vegetative stage, grown on nutrient solid medium and cross-linked their chromatin proteins to DNA by formaldehyde fixation. We isolated chromatin and sonicated it to produce DNA fragments of 300–1,500 bp. We then immunoprecipitated chromatin using antibodies against acetylated histone H3 recognizing acetylated Lys9 and Lys14 residues (#06-599 from Upstate Biotechnology) or the antibody to acetylated histone H4 recognizing acetylated Lys5, Lys8, Lys12 and Lys16 residues (#06-866 from Upstate Biotechnology). We then reversed the cross-links by incubation at 65  $^{\circ}$ C for 5 h. We purified DNA with QIAquick spin columns (QIAGEN) according to the manufacturer's instructions and eluted it in 40  $\mu$ l of 10 mM Tris-HCl, 1 mM EDTA (pH 8.0).

We carried out semiquantitative PCR to amplify the ten different fragments of the *FLC* flowering gene<sup>3,4</sup> (details and primer sequences available on request). All PCR reactions were done in a final volume of 20  $\mu$ l using 1  $\mu$ l of eluted DNA and applying 23–30 cycles (optimized depending on the fragment to avoid saturation of the reaction) of 94  $^{\circ}$ C (30 s), 56  $^{\circ}$ C (45 s) and 72  $^{\circ}$ C (20 s). The amount of amplified DNA was first quantified by image analysis of the DNA bands on agarose gel (visible after 26–30 PCR cycles) using MCID Analysis software (Imaging Research). The results of these quantifications were verified by phosphorimager analysis after hybridization of *FLC* genomic and *ACTIN2* probes on Southern blots containing DNA amplified with 23 cycles (not visible on agarose gels). Data provided in the text are from the first quantification. For each fragment in each genotype sample, the amount of immunoprecipitated DNA was estimated as the amount of amplified DNA normalized to the amount of *ACTIN2* DNA amplified<sup>30</sup>. To estimate the relative amount of immunoprecipitated DNA in the mutants versus the corresponding wild type, we determined the normalized amount of immunoprecipitated DNA in the mutant relative to that of the wild type and calculated the average of the relative value of the ten *FLC* fragments. We carried out, quantified and averaged three independent experiments including the mutant and wild-type genotypes.

**GenBank accession numbers.** *A. thaliana Ler FVE* genomic sequence, AF498101; *A. thaliana Ler FVE* cDNA sequence, AF498102.

*Note: Supplementary information is available on the Nature Genetics website.*

## ACKNOWLEDGMENTS

We thank M. Koornneef, F. Schomburg, R. Amasino and Y. Komeda for providing the seeds of *fve-1*, *fve-2*, *fve-3* and *fve-4* mutants, respectively; M. Koornneef and W. Soppe for providing a *Ler* genomic library; S. Michaels and R. Amasino for a *FLC* cDNA clone; and C. Gutierrez and J. del Pozo for supplying the ZmRBR1 protein and the corresponding antibody. This work has been supported by grants from the European Union and from the Spanish Ministerio de Ciencia y Tecnología. I.A. was supported by predoctoral fellowship from the Instituto Nacional de Investigación y Tecnología Agraria y Alimentaria. C.A.-B. was supported by a contract "Ramón y Cajal" from the Spanish Ministerio de Ciencia y Tecnología.

## COMPETING INTERESTS STATEMENT

The authors declare that they have no competing financial interests.

Received 24 October; accepted 31 December 2003

Published online at <http://www.nature.com/naturegenetics/>

- Simpson, G.G. & Dean, C. Arabidopsis, the rosetta stone of flowering time?. *Science* **296**, 285–289 (2002).
- Koornneef, M., Alonso-Blanco, C., Peeters, A.J.M. & Soppe, W. Genetic control of flowering time in *Arabidopsis*. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **49**, 345–370 (1998).
- Michaels, S.D. & Amasino, R.M. *FLOWERING LOCUS C* encodes a novel MADS-domain protein that acts as a repressor of flowering. *Plant Cell* **11**, 949–956 (1999).
- Sheldon, C.C. *et al.* The *FLF* MADS box gene: a repressor of flowering in *Arabidopsis* regulated by vernalization and methylation. *Plant Cell* **11**, 445–458 (1999).
- Johanson, U. *et al.* Molecular analysis of *FRIGIDA*, a major determinant of natural variation in *Arabidopsis* flowering time. *Science* **290**, 344–347 (2000).
- Koornneef, M., Hanhart, C.J. & Van der Veen, J.H. A genetic and physiological analysis of late flowering mutants in *Arabidopsis thaliana*. *Mol. Gen. Genet.* **229**, 57–66 (1991).
- Sanda, S.L. & Amasino, R.M. Interaction of *FLC* and late-flowering mutations in *Arabidopsis thaliana*. *Mol. Gen. Genet.* **251**, 69–74 (1996).
- Michaels, S.D. & Amasino, R.M. Loss of *FLOWERING LOCUS C* activity eliminates the late-flowering phenotype of *FRIGIDA* and autonomous-pathway mutations, but not responsiveness to vernalization. *Plant Cell* **13**, 935–941 (2001).
- Macknight, R. *et al.* *FCA*, a gene controlling flowering time in *Arabidopsis*, encodes a protein containing RNA-binding domains. *Cell* **89**, 1–20 (1997).
- Schomburg, F.M., Patton, D.A., Meinke, D.W. & Amasino, R.M. *FPA*, a gene involved in floral induction in *Arabidopsis*, encodes a protein containing RNA-recognition motifs. *Plant Cell* **13**, 1427–1436 (2001).
- Simpson, G.G., Dijkwel, P.P., Quesada, V., Henderson, I. & Dean, C. *FY* is an RNA 3' end-processing factor that interacts with *FCA* to control the *Arabidopsis* floral transition. *Cell* **113**, 777–787 (2003).
- Lee, I. *et al.* Isolation of *LUMINIDEPENDENS*: A gene involved in the control of flowering time in *Arabidopsis*. *Plant Cell* **6**, 75–83 (1994).
- Martinez-Zapater, J.M. & Somerville, C.R. Effect of light quality and vernalization on late-flowering mutant of *Arabidopsis thaliana*. *Plant Physiol.* **92**, 770–776 (1990).
- Martinez-Zapater, J.M., Jarillo, J.A., Cruz-Alvarez, M., Roldan, M. & Salinas, J. *Arabidopsis* late-flowering *fve* mutants are affected in both vegetative and reproductive development. *Plant J.* **7**, 543–551 (1995).
- Kenzior, A.L. & Folk, W.R. AtMSI4 and RbAp48 WD-40 repeat proteins bind metal ions. *FEBS Lett.* **440**, 425–429 (1998).
- Li, D. & Roberts, R. WD-repeat proteins: structure characteristics, biological function, and their involvement in human diseases. *Cell. Mol. Life Sci.* **58**, 2085–2097 (2001).
- Hicks, G.R. & Raikhel, N.V. Protein import into the nucleus: an integrated view. *Ann. Rev. Cell Dev. Biol.* **11**, 155–188 (1995).
- Ach, R.A., Taranto, P. & Grussem, W. A conserved family of WD-40 proteins binds to the retinoblastoma protein in both plants and animals. *Plant Cell* **9**, 1595–1606 (1997).
- The *Arabidopsis* genome initiative. Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature* **408**, 796–815 (2000).
- Hennig, L., Taranto, P., Walsler, M., Schönrock, N. & Grussem, W. *Arabidopsis MSI1* is required for epigenetic maintenance of reproductive development. *Development* **130**, 2555–2565 (2003).
- Köhler, C. *et al.* *Arabidopsis* MSI1 is a component of the MEA/FIE Polycomb group complex and required for seed development. *EMBO J.* **22**, 4804–4814 (2003).
- Kaya, H. *et al.* *FASCIATA* genes for chromatin assembly factor-1 in *Arabidopsis* maintain the cellular organization of apical meristems. *Cell* **104**, 131–142 (2001).
- Rossi, V. *et al.* The maize WD-repeat gene *ZmRbAp1* encodes a member of the MSI/RbAp sub-family and is differentially expressed during endosperm development. *Mol. Genet. Genomics* **265**, 576–584 (2001).
- Williams, L. & Graf, G. The retinoblastoma protein - a bridge to heterochromatin. *Trends Plant Sci.* **5**, 239–240 (2000).
- Rossi, V. *et al.* A maize histone deacetylase and retinoblastoma-related protein physically interact and cooperate in repressing gene transcription. *Plant Mol. Biol.* **51**, 401–413 (2003).
- He, Y., Michaels, S.D. & Amasino, R.M. Regulation of flowering time by histone acetylation in *Arabidopsis*. *Science* **302**, 1751–1754 (2003).
- Lee, I., Michaels, S.D., Masshardt, A.S. & Amasino, R.M. The late-flowering phenotype of *FRIGIDA* and mutations in *LUMINIDEPENDENS* is suppressed in the Landsberg *erecta* strain of *Arabidopsis*. *Plant J.* **6**, 903–909 (1994).
- Gray, W.M. *et al.* Identification of an SCF ubiquitin-ligase complex required for auxin response in *Arabidopsis thaliana*. *Genes Dev.* **13**, 1678–1691 (1999).
- Boniotti, M.B. & Gutierrez, C. A cell-cycle-regulated kinase activity phosphorylates plant retinoblastoma protein and contains, in *Arabidopsis*, a CDKA/cyclin D complex. *Plant J.* **28**, 341–350 (2001).
- Johnson, L.M., Cao, X. & Jacobsen, S.E. Interplay between two epigenetic marks: DNA methylation and histone H3 lysine 9 methylation. *Curr. Biol.* **12**, 1360–1367 (2002).