



## Convergence rate of Padé-type approximants for Stieltjes functions

M. Bello Hernández<sup>a,\*</sup>, F. Cala Rodríguez<sup>b</sup>, J.J. Guadalupe<sup>a,1</sup>, G. López Lagomasino<sup>c,2</sup>

<sup>a</sup> *Dpto. de Matemáticas y Computación, Univ. de La Rioja, Logroño, Spain*

<sup>b</sup> *Dpto. de Análisis Matemático, Univ. de La Laguna, Tenerife, Spain*

<sup>c</sup> *Dpto. de Matemáticas, Univ. Carlos III de Madrid, Madrid, Spain*

Received 30 October 1997; received in revised form 20 March 1998

### Abstract

For a wide class of Stieltjes functions we estimate the rate of convergence of Padé-type approximants when the number of fixed poles represents a fixed proportion with respect to the order of the rational approximant. © 1998 Elsevier Science B.V. All rights reserved.

*Keywords:* Orthogonal polynomials; Padé-type approximation

### 1. Introduction

Let  $\gamma > 1$ , by  $f_\gamma$  we denote a continuous almost everywhere positive function on the real line such that

$$\lim_{|x| \rightarrow \infty} f_\gamma(x)|x|^{-\gamma} = 1. \quad (1)$$

In [9], Rakhmanov studied the asymptotic behavior of the sequence  $h_n(d\rho_\gamma; \cdot)$  of orthonormal polynomials with respect to

$$d\rho_\gamma(x) = \exp\{-f_\gamma(x)\} dx, \quad x \in \mathbb{R}. \quad (2)$$

(Within this class of measures, of particular interest are the so-called Freud weights

$$dw_\gamma(x) = \exp\{-|x|^\gamma\} dx$$

\* Corresponding author.

<sup>1</sup> This research was partially supported by DGES under grant PB96-0120-C03-02.

<sup>2</sup> This research was partially supported by DGES under grant PB96-0120-C03-01.

and their orthogonal polynomials.) He proved that

$$\lim_{n \rightarrow \infty} \frac{\log |h_n(d\rho_\gamma; z)|}{n^{1-\gamma^{-1}}} = D(\gamma)|\operatorname{Im} z|, \quad (3)$$

where this limit is uniform on compact subsets of  $\mathbb{C} \setminus \mathbb{R}$ ,

$$D(\gamma) = \frac{\gamma}{\gamma - 1} \left[ \frac{\Gamma((\gamma + 1)/2)}{\Gamma(\gamma/2)} \right]^{(1/\gamma)},$$

and  $\Gamma(\cdot)$  denotes the Gamma function. Set

$$\hat{\rho}_\gamma(z) = \int \frac{d\rho_\gamma(x)}{z - x}.$$

Let  $\pi_n$  denote the  $n$ th diagonal Padé approximant with respect to  $\hat{\rho}_\gamma$ . From Rakhmanov's result it is not hard to deduce that

$$\lim_{n \rightarrow \infty} \frac{\log |\hat{\rho}_\gamma(z) - \pi_n(z)|}{n^{1-\gamma^{-1}}} \leq -2D(\gamma)|\operatorname{Im} z|, \quad (4)$$

uniformly on compact subsets of  $\mathbb{C} \setminus \mathbb{R}$ . We aim to obtain similar results when instead of Padé approximants, Padé-type approximants are used.

Let  $l_n^2$  be a polynomial of degree  $m(n)$  and  $0 \leq m(n) \leq n$ . We define the  $n$ th Padé-type approximants of  $\hat{\rho}_\gamma$  with fixed poles at the zeros of  $l_n^2$  as the unique rational function

$$r_n = \frac{p_n}{q_n l_n^2},$$

where  $p_n$  and  $q_n$  are polynomials which satisfy

- $\deg p_n \leq n - 1$ ,  $\deg q_n \leq n - m(n)$ ,  $q_n \not\equiv 0$ ,
- $(q_n l_n^2 \hat{\rho}_\gamma - p_n)(z) = O(1/(z^{n-m(n)+1}))$ , as  $z = ix \rightarrow \infty$ ,  $x > 0$ .

It is easy to prove (see, e.g., [4]) that

$$0 = \int x^v q_n(x) l_n^2(x) d\rho_\gamma(x), \quad v = 0, \dots, n - m(n) - 1, \quad (5)$$

$$(\hat{\rho}_\gamma - r_n)(z) = \frac{1}{(q_n l_n^2)^2(z)} \int \frac{(q_n l_n^2)^2(x) d\rho_\gamma(x)}{z - x}. \quad (6)$$

If  $m(n) = 0$ , then  $r_n$  is the  $n$ th diagonal Padé approximant with respect to  $\hat{\rho}_\gamma$ . If  $m(n) = n$ , all the poles of the rational approximant are fixed.

In recent years (see, e.g., [1–7]), the rate of convergence of Padé-type and multipoint Padé-type approximants has been studied when the measure defining the function has compact support. We will show that results of type (4) take place for Padé-type approximants when the support of the measure is unbounded. To this end, we will restrict the type of polynomials which carry as their zeros the fixed poles of the Padé-type approximants. In the sequel,  $l_n$  denotes the orthonormal polynomial of degree  $m(n)/2$  with respect to the Freud measure  $dw_\beta(x)$  introduced above. Unless otherwise stated, we take  $\gamma > \beta > 1$ .

We prove

**Theorem 1.** Let  $l_n$  denote the orthonormal polynomial of degree  $m(n)/2$  with respect to the Freud measure  $dw_\beta(x)$  where  $1 < \beta < \gamma$ . Let  $r_n$  denote the  $n$ th Padé-type approximant of  $\hat{\rho}_\gamma$  with fixed poles at the zeros of  $l_n^2$  and assume that

$$\lim_{n \rightarrow \infty} \frac{m(n)}{n} = \theta \in [0, 1).$$

Then

$$\limsup_{n \rightarrow \infty} \frac{\log |\hat{\rho}_\gamma(z) - r_n(z)|}{n^{1-\gamma^{-1}}} \leq -2(1 - \theta)^{1-\gamma^{-1}} D(\gamma) |\operatorname{Im} z|, \tag{7}$$

where convergence takes place uniformly on each compact subset of  $\mathbb{C} \setminus \mathbb{R}$ .

The paper is divided as follows. In Section 2, we give some auxiliary results. Section 3 is devoted to the proof of the theorem stated above and some comments.

## 2. Auxiliary results

Let  $d\rho$  be a finite positive Borel measure on  $\mathbb{R}$ , with an infinite number of points in its support and finite moments. Denote

$$K_j(d\rho, z) = \sup_{p \in \Pi_j, p \neq 0} \frac{|p^2(z)|}{\int |p^2(x)| d\rho(x)}, \tag{8}$$

where  $\Pi_j$  is the set of all polynomials of degree  $\leq j$ .

If  $d\rho = l_n^2 d\rho_\gamma$ , we denote

$$K_{n,j}(z) = K_j(d\rho, z).$$

**Lemma 2.1.** There exist constants  $D > 0$  and  $\alpha \in \mathbb{R}$  such that

$$Dn^\alpha K_j(d\tilde{\rho}_\gamma, z) \leq K_{n,j}(z) \leq K_j(l_n^2 d\rho_\gamma|_{(-n^{1/\gamma}, n^{1/\gamma})}, z), \tag{9}$$

where  $d\tilde{\rho}_\gamma(x) = \exp\{-(f_\gamma(x) - |x|^\beta)\} dx$ ,  $1 < \beta < \gamma$ , and  $l_n^2 d\rho_\gamma|_{(-n^{1/\gamma}, n^{1/\gamma})}$  is the restriction of the measure  $l_n^2 d\rho_\gamma$  to  $(-n^{1/\gamma}, n^{1/\gamma})$ .

**Proof.** From (8) the inequality on the right side of (9) follows directly. On the other hand from Corollary 1.4 in [7] there exist constants  $D_1 > 0$  and  $\alpha_1 \in \mathbb{R}$  such that

$$l_n^2(x) \exp\{-|x|^\beta\} \leq D_1(m(n) + 1)^{\alpha_1}, \quad x \in \mathbb{R}.$$

Since  $0 \leq m(n) \leq n$ , we obtain

$$l_n^2(x) \exp(-|x|^\beta) \leq D_2 n^{\alpha_1}.$$

Thus, if  $p \in \Pi_j$  and  $p \neq 0$ , we get

$$\frac{|p^2(z)|}{\int |p^2(x)| l_n^2(x) d\rho_\gamma(x)} \geq \frac{|p^2(z)|}{D_2 n^{2\alpha} \int |p^2(x)| d\tilde{\rho}_\gamma(x)}$$

and the proof is concluded.  $\square$

**Lemma 2.2.** *Let  $K$  be a compact subset of  $\mathbb{C} \setminus \mathbb{R}$ ,  $1 < \beta < \gamma$ , and  $\lim(m(n)/n) = \theta$ . Then*

$$\liminf_{n \rightarrow \infty} \frac{\log |q_n|(z)}{n^{1-1/\gamma}} \geq (1 - \theta)^{1-1/\gamma} D(\gamma) |\operatorname{Im} z|$$

uniformly on  $K$ , where  $q_n$  is the  $(n - m(n))$ th orthonormal polynomial with respect to  $l_n^2 d\rho_\gamma$  and  $l_n$  denotes the orthonormal polynomial of degree  $m(n)/2$  with respect to the Freud measure  $dw_\beta(x)$ .

**Proof.** Let  $t_k$  be the  $k$ th orthonormal polynomial with respect to  $d\rho$ . From the general theory of orthogonal polynomials, we know (see [5]) that

$$K_j(d\rho, z) = \sum_{k=0}^j |t_k(z)|^2 \geq |t_j(z)|^2, \quad z \in \mathbb{C}, \quad (10)$$

$$K_{j-1}(d\rho, z) = \frac{\tau_{j-1} t_j(z) t_{j-1}(\bar{z}) - t_j(\bar{z}) t_{j-1}(z)}{z - \bar{z}},$$

where  $z \in \mathbb{C} \setminus \mathbb{R}$  and  $\tau_j$  is the leading coefficient of  $t_j$ . Thus, with the aid of (10), we obtain

$$\begin{aligned} K_{j-1}(d\rho, z) &= \frac{\tau_{j-1} \operatorname{Im}(t_j(z) \overline{t_{j-1}(z)})}{\tau_j \operatorname{Im} z} \\ &\leq \frac{\tau_{j-1} |t_j t_{j-1}(z)|}{\tau_j |\operatorname{Im} z|} \\ &\leq \frac{\tau_{j-1} |t_j(z)| K_{j-1}^{1/2}(z)}{\tau_j |\operatorname{Im} z|}. \end{aligned}$$

This inequality yields

$$K_{j-1}(d\rho, z) \leq \frac{\tau_{j-1}^2 |t_j(z)|^2}{\tau_j^2 |\operatorname{Im} z|^2}, \quad (11)$$

therefore,

$$K_j(d\rho, z) = K_{j-1} + |t_j(z)|^2 \leq \left[ \frac{\tau_{j-1}^2}{\tau_j^2 |\operatorname{Im}(z)|^2} + 1 \right] |t_j(z)|^2. \quad (12)$$

On the other hand,

$$\begin{aligned} \frac{1}{\tau_j^2} &= \inf_{p=z/\dots} \int |P^2(x)| \, d\rho(x) \\ &\leq \int \left| x \frac{t_{j-1}(x)}{\tau_{j-1}} \right|^2 \, d\rho(x), \end{aligned}$$

or what is the same

$$\frac{\tau_{j-1}^2}{\tau_j^2} \leq \int |xt_{j-1}|^2 \, d\rho(x).$$

If  $d\rho(x)$  satisfies (2), there exist constants  $D_1, D_2, D_3 > 0$  such that for all  $k \in \mathbb{N}$  and  $p \in \Pi_k$ , we have (see Theorem 2.6 in [8])

$$\int |p^2(x)| \, d\rho(x) \leq D_2 \int_{-D_1 k^{1/\gamma}}^{D_1 k^{1/\gamma}} |p(x)|^2 \, d\rho(x),$$

in particular,

$$\begin{aligned} \frac{\tau_{j-1}^2}{\tau_j^2} &\leq D_2 \int_{-D_1 j^{1/\gamma}}^{D_1 j^{1/\gamma}} |xt_{j-1}(x)|^2 \, d\rho(x), \\ &\leq D_3 j^{2/\gamma}. \end{aligned} \tag{13}$$

Take  $d\rho(x) = d\tilde{\rho}_\gamma(x) = \exp\{-(f_j(x) - |x|^\beta)\} \, dx$ . Since  $1 < \beta < \gamma$ , the function  $f_j(x) - |x|^\beta$  satisfies (1). Using (3), (10), (12), and (13) one obtains

$$\lim_{n \rightarrow \infty} \frac{\log K_n(d\tilde{\rho}_\gamma, z)}{n^{1-1/\gamma}} = 2D(\gamma)|\text{Im}(z)|. \tag{14}$$

This result appears in [9], Lemma 4.

For  $d\rho(x) = l_n^2(x) \, d\rho_\gamma(x)$  and  $j = n - m(n)$ , (12) gives

$$K_{n, n-m(n)}(z) \leq \left[ \frac{\tau_{n, n-m(n)-1}^2}{\tau_{n, n-m(n)}^2} + 1 \right] |q_n(z)|^2, \tag{15}$$

where  $q_n$  is the  $(n - m(n))$ th orthonormal polynomial with respect to  $|l_n|^2 \, d\rho_\gamma$  and  $\tau_{n, n-m(n)}$  its leading coefficient. Notice that, infinite–finite range  $L_2$  estimates give as above

$$\frac{\tau_{n-m(n)-1}^2}{\tau_{n-m(n)}^2} \leq D_4 n^{2/\gamma}. \tag{16}$$

From the first inequality in (9), (14)–(16), we obtain

$$\begin{aligned} \liminf_{n \rightarrow \infty} \frac{\log |q_n(z)|}{n^{1-1/\gamma}} &\geq \lim_n \left( \frac{n - m(n)}{n} \right)^{1-1/\gamma} \frac{\log |K_{n-m(n)}(d\tilde{\rho}_\gamma, z)|}{2(n - m(n))^{1-1/\gamma}} \\ &= (1 - \theta)^{1-1/\gamma} D(\gamma)|\text{Im}(z)| \end{aligned}$$

and the proof is finished.  $\square$

### 3. Proof of Theorem 1

Let  $K$  be a compact subset of  $\mathbb{C} \setminus \mathbb{R}$ , then there exists  $D_1 = D_1(K) > 0$  such that

$$|z - x| \geq D_1, \quad z \in K, \quad x \in \mathbb{R}.$$

Using (6) and the orthonormality of  $q_n$ , we get

$$\begin{aligned} |(\hat{\rho}_\gamma - r_n)(z)| &= \left| \frac{1}{(q_n l_n)^2(z)} \int \frac{(q_n l_n)^2(x)}{z - x} d\rho_\gamma \right| \\ &\leq \frac{1}{D_1 |(q_n l_n)^2(z)|}. \end{aligned}$$

Now, from Lemma 2.2 and (3) as applied to the sequence  $\{l_n\}$ , we obtain (7).  $\square$

**Corollary 1.** *Under the assumptions of Theorem 1*

$$r_n \rightarrow \hat{\rho}_\gamma,$$

*uniformly on each compact set of  $\mathbb{C} \setminus \mathbb{R}$ .*

**Proof.** It is immediate from the fact that the right-hand side of (7) is continuous and negative on  $\mathbb{C} \setminus \mathbb{R}$ .  $\square$

**Remark 1.** In the case when  $\theta = 1$  and  $1 < \beta = \gamma$  it is possible to construct examples where there is divergence. For example, taking  $m(n) = n$  and  $f_\gamma(x) = |x|^\gamma - |x|^{\gamma'}$  with  $\gamma' < \gamma$  sufficiently close to  $\gamma$ . For this reason we do not discuss this limiting situation.

**Remark 2.** When  $1 < \gamma < \beta$  and  $m(n) = n$  there is always divergence.

### References

- [1] A. Ambroladze, H. Wallin, Padé type approximants of Markov and meromorphic functions, *J. Approx. Theory* 88 (1997) 354–369.
- [2] A. Ambroladze, H. Wallin, Convergence of rational interpolants with preassigned poles, *J. Approx. Theory* 89 (1997) 238–256.
- [3] F. Cala, G. López, From Padé to Padé-type approximants. Exact rate of convergence, in: M. Florenzano et al. (Eds.), *Proc. 2nd Internat. Conf. on Approx. and Opt. in the Caribbean*, Peter Series in Approx. and Optimization, vol. 8, 1995, pp. 155–163.
- [4] F. Cala, G. López, Multipoint Padé-type approximants. Exact rate of convergence, *Constr. Approx.* 14 (2) (1998) 259–272.
- [5] G. Freud, *Orthogonal Polynomials*, Pergamon Press, Budapest, 1966.
- [6] A.A. Gonchar, On convergence of Padé approximants for some classes of meromorphic functions, *Mat. Sb.* 97 (139) (1975) 4; English Transl. in: *Math. USSR Sb.* 26 (1975) 555–575.

- [7] A.L. Levin, D.S. Lubinsky, Christoffel functions, Orthogonal Polynomials, and Nevai's Conjecture for Freud Weights, *Constr. Approx.* 8 (1992) 463–535.
- [8] H.N. Mhaskar, E.B. Saff, Where does the  $L_p$  norm of a weighted polynomial live?, *Trans. Amer. Math. Soc.* 303 (1987) 109–124.
- [9] E.A. Rakhmanov, On the asymptotic properties of polynomials orthogonal on the real axis, *Mat. Sb.* 119 (161) (1982) 161–203; English Transl. in: *Math. USSR Sb.* 47 (1984) 155–193.