

CORRECTION



Correction to: Fetal exposure to phthalates and bisphenols and childhood general and organ fat. A population-based prospective cohort study

Chalana M. Sol, Susana Santos, Liesbeth Duijts, Alexandros G. Asimakopoulos, Maria-Pilar Martinez-Moral, Kurunthachalam Kannan, Elise M. Philips, Leonardo Trasande  and Vincent W. V. Jaddoe 

© The Author(s), under exclusive licence to Springer Nature Limited 2022

International Journal of Obesity (2022) 46:1411–1412; <https://doi.org/10.1038/s41366-022-01126-y>

Correction to: *International Journal of Obesity* <https://doi.org/10.1038/s41366-020-00672-7>, published online 12 September 2020

The original version of this article unfortunately contained a mistake in table 1. On page 2227, in Table 1, the limit of detection in nmol/L for two of the chemicals has been noted incorrectly. The limit of detection (LOD) for mono-2-heptyl phthalate (mHpP) in nmol/L was noted in the table as 0.97 instead of 1.13 and the

limit of detection for Bisphenol F (BPF) in nmol/L was noted in the table as 0.86 instead of 0.90. These typing mistakes did not influence the analysis, results or conclusions, as the authors did not perform any calculations based on the LOD-values as presented in the table, but for all analysis the original values were used, which the authors have verified were correct. The correct table can be found below. The original article has been corrected.

Table 1. Urinary concentrations of phthalates and bisphenols during pregnancy ($n = 1128$).

| | Limit of detection (nmol/L) | First trimester Median (25 th –75 th percentile) | Second trimester Median (25 th –75 th percentile) | Third trimester Median (25 th –75 th percentile) |
|---|-----------------------------|---|--|---|
| Phthalic Acid (PA) (nmol/L) | 6.68 | 340.6 (180.4–724.4) | 899.7 (355.4–1682.3) | 406.9 (203.1–794.3) |
| Low-molecular-weight phthalates (LMWP) (nmol/L) | – | 1 094.8 (434.8–2910.0) | 550.5 (227.2–1441.8) | 991.8 (404.3–2480.7) |
| Monomethylphthalate (mMP) (nmol/L) | 0.33 | 29.8 (15.1–54.8) | 18.9 (9.8–34.7) | 21.8 (10.9–44.9) |
| Monoethylphthalate (mEP) (nmol/L) | 0.31 | 717.7 (208.1–2466.6) | 347.5 (121.1–1101.8) | 642.9 (223.2–2076.9) |
| Mono-isobutylphthalate (mIBP) (nmol/L) | 0.40 | 95.5 (42.7–201.5) | 39.2 (19.9–77.5) | 76.6 (40.8–152.9) |
| Mono-n-butylphthalate (mBP) (nmol/L) | 0.63 | 72.4 (30.4–138.6) | 41.9 (23.3–81.5) | 53.5 (27.4–110.0) |
| High-molecular-weight phthalates (HMWP) (nmol/L) | – | 214.3 (113.1–396.8) | 128.8 (70.6–234.8) | 165.9 (95.2–291.3) |
| Monobenzylphthalate (mBzBP) (nmol/L) | 0.59 | 22.1 (8.8–46.6) | 19.6 (8.3–40.5) | 12.0 (4.0–24.6) |
| Mono-hexylphthalate (mHxP) (nmol/L) | 0.24 | 0.9 (0.3–2.0) | NA | NA |
| Mono-2-heptylphthalate (mHpP) (nmol/L) | 1.13 | 2.1 (0.8–5.3) | NA | NA |
| Monocyclohexyl-phthalate (mCHP) (nmol/L) | 0.16 | 0.1 (0.1–0.1) | NA | NA |
| Di-2-ethylhexylphthalate (DEHP) (nmol/L) | – | 170.1 (88.6–313.8) | 95.8 (51.2–178.3) | 137.7 (77.5–248.1) |
| Mono-(2-ethyl-5-carboxy-pentyl)phthalate (mECPP) (nmol/L) | 0.94 | 52.0 (26.1–99.93) | 33.2 (18.1–62.0) | 58.0 (30.4–108.2) |
| Mono-(2-ethyl-5-hydroxy-hexyl)phthalate (mEHP) (nmol/L) | 0.27 | 40.2 (19.7–77.6) | 18.4 (10.0–36.1) | 34.1 (17.9–66.3) |
| Mono-(2-ethyl-5-oxohexyl)phthalate (mEOHP) (nmol/L) | 0.14 | 26.4 (11.9–52.5) | 24.8 (11.9–54.4) | 24.5 (13.1–47.2) |
| Mono-[(2-carboxymethyl)-hexyl] phthalate (mCMHP) (nmol/L) | 0.13 | 4.5 (24.5–80.8) | 12.6 (7.1–23.1) | 10.8 (5.8–20.4) |
| Di-n-octylphthalate (DNOP) | – | | | |
| Mono(3-carboxypropyl)- phthalate (mCPP) (nmol/L) | 0.03 | 5.7 (3.0–10.7) | 3.5 (2.0–6.6) | 7.1 (3.8–12.4) |
| Bisphenols (nmol/L) | – | 9.05 (3.52–20.3) | 6.3 (2.9–13.6) | 9.1 (4.2–18.8) |
| Bisphenol A (BPA) (nmol/L) | 0.66 | 4.8 (1.0–12.3) | 5.8 (2.6–12.4) | 6.5 (2.7–12.8) |
| Bisphenol S (BPS) (nmol/L) | 0.20 | 0.7 (0.1–2.5) | 0.1 (0.1–0.4) | NA |
| Bisphenol F (BPF) (nmol/L) | 0.90 | 0.6 (0.6–2.1) | NA | 0.6 (0.6–2.5) |

Values represent medians (25th–75th percentiles). Absolute urinary concentration of the limit of detection (in nmol/L urine), grouped exposures (in nmol/L urine), and individual exposures (in nmol/L urine) with concentrations below the limit of detection imputed as limit of detection/square root of 2. Only values that are included in the calculation of the grouped exposures are included in this table. NA not applicable due to >80% of concentrations below limit of detection.