



Genomic Analysis of Bacteriocin-Producing Staphylococci: High Prevalence of Lanthipeptides and the Micrococcin P1 Biosynthetic Gene Clusters

Rosa Fernández-Fernández¹ · Ahmed M. A. Elsherbini^{2,3,4} · Carmen Lozano¹ · Agustí Martínez¹ · María de Toro⁵ · Myriam Zarazaga¹ · Andreas Peschel^{2,3,4} · Bernhard Krismer^{2,3,4} · Carmen Torres¹

Accepted: 30 June 2023
© The Author(s) 2023

Abstract

Bacteriocins are antimicrobial peptides produced by bacteria. This study aimed to in silico analyze the presence of bacteriocin gene clusters (BGCs) among the genomes of 22 commensal *Staphylococcus* isolates from different origins (environment/human/food/pet/wild animals) previously identified as bacteriocin producers. The resistome and plasmidome were studied in all isolates. Five types of BGC were detected in 18 genomes of the 22 bacteriocin-producing staphylococci included in this study: class I (Lanthipeptides), class II, circular bacteriocins, the non-ribosomal-peptide lugdunin and the thiopeptide micrococcin P1 (MP1). A high frequency of lanthipeptides was detected in this collection: BGC variants of BSA, bacCH91, and epilancin15X were identified in two *Staphylococcus aureus* and one *Staphylococcus warneri* isolates from food and wild animals. Moreover, two potentially new lanthipeptide-like BGCs with no identity to database entries were found in *Staphylococcus epidermidis* and *Staphylococcus simulans* from food and wild animal, respectively. Interestingly, four isolates (one *S. aureus* and one *Staphylococcus hominis*, environmental origin; two *Staphylococcus sciuri*, food) carried the MP1 BGC with differences to those previously described. On the other hand, seven of the 22 genomes (~32%) lacked known genes related with antibiotic or disinfectant-acquired resistance mechanisms. Moreover, the potential carriage of plasmids was evaluated, and several Rep-proteins were identified (~73% of strains). In conclusion, a wide variety of BGCs has been observed among the 22 genomes, and an interesting relationship between related *Staphylococcus* species and the type of bacteriocin has been revealed. Therefore, bacteriocin-producing *Staphylococcus* and especially coagulase-negative staphylococci (CoNS) can be considered good candidates as a source of novel bacteriocins.

Keywords Bacteriocins · BGC · CoNS · Lanthipeptides · Micrococcin P1

Rosa Fernández-Fernández and Ahmed M. A. Elsherbini contributed equally to this work.

✉ Carmen Torres
carmen.torres@unirioja.es

- ¹ Area of Biochemistry and Molecular Biology, OneHealth-UR Research Group, University of La Rioja, 26006 Logroño, Spain
- ² Department of Infection Biology, Interfaculty Institute of Microbiology and Infection Medicine, University of Tübingen, 72076 Tübingen, Germany
- ³ Cluster of Excellence EXC 2124 Controlling Microbes to Fight Infections, Tübingen, Germany
- ⁴ German Center for Infection Research (DZIF), Partner Site Tübingen, Tübingen, Germany
- ⁵ Genomics and Bioinformatics Core Facility, Center for Biomedical Research of La Rioja, Logroño, Spain

Introduction

Antimicrobial resistance (AMR) represents one of the biggest challenges of modern medicine worldwide [1]. This AMR global problem requires novel antimicrobial agents and strategies to overcome the threat by various pathogens, including multidrug-resistant (MDR) and zoonotic bacteria [2, 3]. About 35% of all drugs and 60–80% of all antimicrobial products have originated from natural products, and the development of innovative products derived from natural substances is gaining attention for pharmaceutical and therapeutic uses [4–6]. However, financial pressure from drug companies as well as difficulties in the isolation and identification of new natural compounds have severely limited the discovery rate from these important sources [5, 6].

Typically, bacteriocins are ribosomally synthesized peptides, encoded by operons including the structural genes

whose products are often post-translationally modified by specific enzymes encoded by adjacent genes. In these genetic systems, regulatory genes, immunity genes, transporters, and other genes that encode accessory proteins are also found [7–9]. Traditionally, the identification of novel bacteriocins has used cultivation-based approaches that involved screening of numerous isolates for antimicrobial activity, followed by long-term biochemical characterization [10, 11]. More recently, genome mining has become an important methodology in the discovery of novel natural products with antimicrobial activity [7]. In this respect, ribosomally synthesized and post-translationally modified peptides (RiPPs) of bacterial origin are natural compounds that are highly attractive candidates for antibacterial prevention and therapy [7, 12]. Nevertheless, other types of bacteriocins can be also very relevant such as Aureocins or Epidermicin NI01 [13].

Bacteriocins have been divided in three major classes depending on their amino acid composition, chemical structure, complexity, mode of action, and steps involved in their production (synthesis, transport, and immunity): class I, small post-translationally modified peptides; class II, unmodified bacteriocins; and class III, larger and thermo-labile peptides [14, 15]. Moreover, recent reviews have proposed the inclusion of additional classes among Gram-positive bacteriocins: class IV (circular), class V (sactipeptides), class VI (thiopeptides) and non-ribosomal peptides (NRP) [13, 14, 16–22].

Within the last few years, increasing numbers of bacteriocins have been isolated and identified from Gram-positive microorganisms [15], and the lanthipeptides (class I) belong to the most frequently found [23]. These peptides represent a promising type of natural antibacterial molecules, active against many Gram-positive pathogens, including antibiotic-resistant strains such as methicillin-resistant *S. aureus* (MRSA) and vancomycin-resistant enterococci (VRE) [24].

Environmental members of the phylum Actinobacteria and Firmicutes including bacilli and lactic acid bacteria, and many Proteobacteria such as *Escherichia coli* are well-known as producers of biologically active substances [25–27]. In addition, *Staphylococcus* species, in particular coagulase-negative staphylococci (CoNS) isolated from a wide variety of natural sources, have also been found to be interesting sources of BGCs for bacteriocins [28–30].

In previous studies carried out by our group [31–33], a collection of 92 bacteriocin-producing staphylococci was detected in a screening approach performed with 1205 staphylococci of different species and origins (~7.7% were bacteriocin producers) (Table S1). Twenty-two of these isolates of 11 species, both coagulase-positive and negative staphylococci (CoPS and CoNS, respectively), were included in this study for further characterization. Isolates were selected under a OneHealth perspective based on their antimicrobial activity profile (high production) and their origin. The specific objectives of this study

were (1) to predict and analyze the presence of BGCs in the genomes of the 22 selected bacteriocin-producing *Staphylococcus* isolates and (2) to investigate the novelty or the homology between our BGCs after comparison with those previously reported in databases.

Material and Methods

Isolates Included in the Study

The genomes of 22 *Staphylococcus* isolates, both CoPS and CoNS, of eleven different species and six origins, were included in this study for genomic comparison. The bacteriocin-producing isolates were as follows (number of isolates, identification code): (a) *S. aureus* from food ($n=1$, X3410), environment ($n=1$, C5802), and wild mammal ($n=1$, C8609); (b) *Staphylococcus pseudintermedius* from dogs ($n=2$, C4502, C8478) and human ($n=1$, C8189); (c) *Staphylococcus chromogenes* from wild mammals ($n=2$, C9838, C9727); (d) *Staphylococcus hyicus* from wild mammals ($n=2$, C9581, C9585); (e) *S. sciuri* recently reclassified as *Mammaliococcus sciuri* [34, 35] were recovered from wild birds ($n=3$, C9213, C9179, C9529) and food ($n=2$, X3011, X3041); (f) *Staphylococcus lugdunensis* from humans ($n=2$, C9161, C9954); (g) *S. hominis* of environmental origin ($n=1$, C5835); (h) *S. warneri* ($n=1$, X2969) from food; (i) *S. simulans* recovered from a wild mammal ($n=1$, C9832); (j) *S. epidermidis* ($n=1$, X3009), from food and (k) *Staphylococcus xylosus* ($n=1$, C9255) isolated from wild birds. The characteristics of the 22 isolates analyzed in this study are indicated in Table S2.

DNA Extraction, Amplification, and Sequencing

The NucleoSpin microbial DNA Kit (Macherey and Nagel, Germany) was used for DNA extraction. For it, 2 ml of a 24-h grown culture in normal BM medium at 37 °C were pelleted (40 mg) by centrifugation (10 min at 10,000 rpm), and the FastPrep-24 Classic homogenizer (MP Biomedicals) was used for mechanical disruption. Genomic DNA was subsequently purified according to the manufacturers' instructions.

The genomic libraries for eight *Staphylococcus* isolates (X2969, X3009, X3041, C8609, C9832, C9581, C9585, and C9838) were prepared with the TruSeq DNA PCR-Free Kit (Illumina) by the Genomics and Bioinformatics Core Facility (CIBIR), whereas the genomic libraries of the other isolates (C4502, C9161, C9179, C9213, C9255, C9529, C9727, C9954, X3011, and X3410) were prepared with the Nextera XT library preparation kit (Illumina) by the NGS Competence Center Tübingen (NCCT). In both cases, libraries were sequenced in different runs on an Illumina MiSeq platform with the MiSeq Reagent Kit v3 (Illumina, 150 cycle). The

genomes of the isolates C5802, C5835, C8478, and C8189 were previously obtained in collaborative studies of the OneHealth-UR research group and all isolates belong to the strain collection of the University of La Rioja.

The Whole Genome project has been deposited at GenBank under the accession PRJNA974190 and the respective accession numbers are included in Table S2.

Data Analysis

Raw reads quality was checked using FastQC (v0.11.8) [36]. Trimming of raw reads, de novo assembly, and assembly polishing were performed with Trimmomatic (v0.39) [37], SPAdes (v3.15.5), and Pilon (v1.24), respectively [38, 39], through the Shovill pipeline [40]. Then, assembled genomes were functionally annotated using Prokka (v1.14.6) [41].

Using protein files exported from the Prokka annotations in FASTA format, a maximum likelihood tree was computed via the RaxML program (v8.2.12) [42], applied inside PhyloPhlAn (v3.0.60) pipeline using the PhyloPhlAn protein markers database and with the default diversity parameter for genus/family level phylogenies [43]. The tree was visualized using the ggtree package (v3.4.4) of R (v4.2.1) [44, 45].

Resistome and Plasmidome

The resistance genotypes and the presence of plasmid replication (*rep*) genes were studied through ResFinder and PlasmidFinder web tools, respectively [46]. In all cases, the default parameters with relaxed mode were used and *S. aureus* was selected as reference species.

Bacteriocin Gene Clusters Prediction

To predict and analyze homologous bacteriocin gene clusters (BGCs) in the 22 genomes, all annotated genomes in GenBank format were uploaded to antiSMASH (v6.1.1) [47] and BAGEL4 [48] with all parameters and a relaxed mode. For BGC prediction, the genomes were considered as a pool of contigs both including chromosome and plasmid regions.

To extract homologous BGCs of micrococcin P1 (MP1) from *Macrococcus caseolyticus*, pBac115 plasmid sequence (accession number KM613043.1) was used as reference, and the protein sequences identified by antiSMASH were blasted against the NCBI protein database using the Cblaster (v1.3.15) tool with default parameters [49]. To simplify the output, unique non-redundant homologous BGCs per species were aligned against the *M. caseolyticus* BGC and the MP1 genomes from this study using the Clinker (v0.0.25) tool with the default parameters [50].

To investigate whether all identified micrococcin P1 BGCs were encoding the identical thiopeptide, the structural protein sequences were extracted from all BGCs of interest. Then, they were mutually aligned by MAFFT (v7.310) and

visualized with Jalview (v2.11.2.0) according to ClustalX color scheme [51].

Moreover, identity comparisons between the MP1 plasmid from *S. hominis* S34-1 (CP040733.1) and the MP1-producing isolates (*S. sciuri* X3041 and X3011, *S. hominis* C5835, *S. aureus* C5802) were performed with Clustal W2 and Emboss Needle [52], and Clinker (v0.0.25) tools [50].

Results

The complete genomes of 22 bacteriocin-producing *Staphylococcus* spp. isolates, both CoPS and CoNS were obtained by WGS and annotation. Phenotypic and molecular characteristics of the isolates as well as results of resistome and plasmidome analyses are presented in Tables S2 and S3. Several bacteriocin-encoding gene clusters were identified, which are summarized in Table 1.

According to the analysis of the 22 genomes included in this study, seven out of 22 (~32%) isolates lacked of known antibiotic or disinfectant resistance-related genes. The remaining 15 isolates (~68%) carried at least one gene associated with antibiotic resistance (beta-lactams, macrolides, lincosamides, streptogramins and aminoglycosides (LSA), tetracyclines, phenicol, fusidic acid, and fosfomycin) or disinfectant agents. Regarding the plasmidome, a wide variety of Rep protein genes were identified in 16 of the studied genomes. Among these genomes, eight carried *rep* and antimicrobial resistance genes and the other eight only carried *rep* genes (Table S2). Interestingly, isolates C9727, C9954, and C9161 did not carry any potential antimicrobial resistance gene, disinfectant-related resistance mechanism or plasmids (lack of Rep protein genes) (Tables S2 and S3).

The use of antiSMASH and BAGEL4 software revealed the presence of a large number and wide diversity of BGCs in the various genomes (Table 1). Comparison of all predicted BGCs identified five types of BGCs in 18 of the 22 genomes of bacteriocin-producing isolates from different species and origins included in this study (Fig. 1). The BGCs from this study were assigned to class I (lanthipeptides), class II (bacSp222, lactococcin972 and blp), circular bacteriocins, NRPs (lugdunin), and thiopeptide (MP1), as follows:

1. Class I: Seven out of the 22 genomes (~32%) carried a lanthipeptide-like BGCs with identities to database entries. The isolates were recovered from wild animals (*S. aureus* C8609 and *S. simulans* C9832) or food (*S. aureus* X3410, *S. warneri* X2969, *S. epidermidis* X3009). Interestingly, two of these isolates (C9832 and X3009) seemed to carry currently unknown lanthipeptide-encoding BGCs. Moreover, a type V lanthipeptide was detected in *S. hyicus* isolate C9585 recovered from a wild animal (wild boar), and strong identity was

Table 1 Details of the type of bacteriocin gene clusters (BGC) predicted both by antiSMASH or BAGEL4 among the 22 genomes included in this study

Isolate	Specie	Origin	BGC type(s) ^a	BGC prediction ^{a,b,c}
X3041	<i>S. sciuri</i>	Food	Class II, thiopeptide	Lactococcin972, MP1
X3011	<i>S. sciuri</i>	Food	Class II, thiopeptide	Lactococcin972, MP1
C9179	<i>S. sciuri</i>	Wild bird	NRPs	None ^c
C9213	<i>S. sciuri</i>	Wild bird	NRPs	None ^c
C9529	<i>S. sciuri</i>	Wild bird	NRPs	None ^c
C9585	<i>S. hyicus</i>	Wild mammal	Lanthipeptide	Lanthipeptide V
C9581	<i>S. hyicus</i>	Wild mammal	RiPP	Circular
C9838	<i>S. chromogenes</i>	Wild mammal	RiPP	2 Circular
C9727	<i>S. chromogenes</i>	Wild mammal	Lanthipeptide	Homincin variant
C9832	<i>S. simulans</i>	Wild mammal	Class II, lanthipeptide	Lactococcin972, new lanthipeptide
C5835	<i>S. hominis</i>	Environmental	Thiopeptide	MP1
C9255	<i>S. xylosum</i>	Wild bird	Class II, RiPP	Lactococcin972, blp, circular
X2969	<i>S. warneri</i>	Food	Lanthipeptide	Epilancin15X variant
X3009	<i>S. epidermidis</i>	Food	Lanthipeptide	New lanthipeptide
C8478	<i>S. pseudintermedius</i>	Pet	Class II	Bacsp222
C8189	<i>S. pseudintermedius</i>	Human	Class II	Bacsp222
C4502	<i>S. pseudintermedius</i>	Pet	NRPs	None ^c
C5802	<i>S. aureus</i>	Environmental	Class II, thiopeptide	Lactococcin972, MP1
C8609	<i>S. aureus</i>	Wild mammal	Class II, lanthipeptide	Lactococcin972, BSA
X3410	<i>S. aureus</i>	Food	Lanthipeptide	BacCH91
C9954	<i>S. lugdunensis</i>	Human	NRPs	Lugdunin
C9161	<i>S. lugdunensis</i>	Human	NRPs	Lugdunin

^aAbbreviation: *MPI* micrococcin P1, *NRPs* non-ribosomal-peptide-synthetase, *RiPP* ribosomally synthesized and post-translationally modified peptide

^bBacteriocin names: bacsp222, BSA, bacCH91

^cNone: in these cases, the relation of the predicted secondary metabolites with antimicrobial substances could not be verified

- found between the BGC predicted in *S. chromogenes* C9727 (wild boar) and the structural gene coding for the lanthipeptide homincin.
- Class II: Genes for lactococcin972-like peptides were detected among 6 out of 22 genomes (~36%) from (a) three isolates recovered from wild animals (*S. aureus* C8609, *S. xylosum* C9255 and *S. simulans* C9832); (b) two *S. sciuri* isolates from food samples (X3041 and X3011); and (c) one *S. aureus* C5802 isolate recovered from an environmental sample. Moreover, part of the bacSp222 bacteriocin gene cluster was detected in two *S. pseudintermedius* (C8478 and C8189) isolates recovered from the same dog–human zoonosis case and a *blp*-like bacteriocin operon was identified in *S. xylosum* C9255.
 - BGCs for circular bacteriocins were identified in three isolates recovered from wild animals (*S. chromogenes* C9838, *S. hyicus* C9581, and *S. xylosum* C9255). The predicted BGCs revealed high identity with uberolysin/circularinA and aureocyclin 4185 described in the databases for other species.

- Non-ribosomal-peptide-synthetase genes (NRPS) coding for lugdunin were detected in two *S. lugdunensis* isolates (C9161 and C9954) recovered from humans.
- Four isolates (~18%) carried a BGC coding for the thiopeptide micrococcin P1: *S. aureus* C5802 and *S. hominis* C5835 from environmental samples and two *S. sciuri* X3011 and X3041 from food.

Class I: Lanthipeptides

The incidence of lanthipeptides was particularly noteworthy and detected in ~32% of the evaluated bacteriocin-producing isolates ($n = 7$) (Fig. 1). A comparison between the lanthipeptide BGC operons was performed following the same criteria in all the used sequences. Figure 2 represents the genetic environment of the BGCs detected in this study (*S. aureus* X3410 and C8609, *S. hyicus* C9585, *S. warneri* X2969, *S. epidermidis* X3009, *S. simulans* C9832 and *S. chromogenes* C9727) and those previously described (accession number): bacCH91 (JQ655767), hyicin3682




Fig. 1 Maximum likelihood tree of protein markers of the 22 bacteriocin-producing isolates of different origins (environmental, food, human, pet, and wild animal both birds and mammals). The bacteriocin gene clusters (BGC) identified after in silico analysis are marked with symbols and colors: class I, lanthipeptides (blue star); class II, lactotoccin972 (purple circle) and bacsp222 (yellow cross); circu-

lar (green square), NRPS (red rhombus), thiopeptide (pink triangle). Abbreviation of specie: SA *S. aureus*, SP *S. pseudintermedius*, SSC *S. sciuri*, SCh *S. chromogenes*, SSi *S. simulans*, SX *S. xylosus*, SL *S. lugdunensis*, SH *S. hominis*, SE *S. epidermidis*, SW *S. warneri*. This figure is in color in the online version but in black and white in the print version

(KY021154), epidermin (X62386), BSA (BA000033.2), pep5 (Z49865), and epilancin15X (JQ979180).

The *S. aureus* X3410 and C8609 BGC revealed a high degree of identity to the bacteriocin bacCH91 (JQ655767), BSA (BA000033.2), epidermin (X62386), and hyicin3682 (KY021154). On the other hand, the *S. warneri* X2969 BGC showed a close identity with the coding operon of

epilancin15X (JQ979180) also related with pep5 BGC (Z49865). Regarding *S. epidermidis* X3009 BGC, slight identities were detected between the predicted structural genes but not with the rest of the encoding genes. Moreover, the *S. simulans* C9832 and *S. hyicus* C9585 lanthipeptide operons were classified separately from the rest of BGCs (Fig. 2).




Fig. 2 Genetic environment comparison between the bacteriocin gene clusters (BGCs) coding for lanthipeptides (class I) in the antimicrobial-producing isolates detected in this study and those selected

as references from the databases. This figure is in color in the online version but in black and white in the print version

Finally, the BGC predicted on the *S. chromogenes* C9727 genome was identified as hominycin. Since only the partial amino acid sequence of this bacteriocin has been reported [53], further studies will be performed to determine their identity.

Comparative analysis presented in Fig. 3 revealed 100% amino acid identity between the bacteriocin structural peptide of *S. aureus* X3410 and C8609 and the BacCH91 structural peptide. Moreover, only slight differences were observed when comparing the bacteriocin structural pre-peptides of *S. aureus*

C8609 and X3410 with the BsaA2 (83% of identity) and BsaA1 (92%) amino acid sequences. In relation to the BGC of *S. warneri* X2969, an 82% of identity was detected with the Epilancin15X structural pre-peptide. In addition, antiSMASH analysis allowed us to identify a lanthipeptide V-like BGC in the *S. hyicus* C9585 genome. Noteworthy, the BGCs of bacteriocin-producing *S. epidermidis* X3009 and *S. simulans* C9832

did not show homologues in the database, indicating the detection of two putatively new lanthipeptides (Fig. 3).

Class II: Bacteriocins

Genome analysis also led to the identification of three types of class II bacteriocins. The BGC for lactococcin972 was




Fig. 3 Sequence alignments of the structural peptides coding for lanthipeptide-like bacteriocins (detected in this study*) and those previously described and published in public databases used as reference. **A** *S. warneri* X2969 and the pre-peptide of epilancin15X and pep5 bacteriocins. **B** Hyicin 3682, epidermin, BsaA1, BsaA2, bacCH91

peptides with those detected in *S. aureus* C8609 isolate and *S. aureus* X3410. **C** Structural peptides detected among the genomes included in this study. * Symbol to indicate the structural peptide of the isolates included in this study. This figure is in color in the online version but in black and white in the print version

predicted in six genomes, and after genomic environment comparisons, all but one potential lactococcin972 BGCs predicted revealed identity with the structural gene of the *L. lactis*_pBL1_NC004955.1 reference sequence (Fig. S1). Moreover, high identities were observed when considering the lactococcin972 BGC predicted among the genomes included in this study, specially between *S. aureus* C8609 and C5802 and *S. sciuri* X3041 and X3011. Contrary, the BGC predicted to code for lactococcin972 in the *S. simulans* C9832 genome did not show relevant identity neither with the reference sequences considered nor with the rest of the predicted operons.

We also detected the identical structural gene of the bacSp222 bacteriocin in the genomes of the two *S. pseudintermedius* isolates recovered from the same human-pet zoonosis case. This bacteriocin was described for the first time in the plasmid p222 of *S. pseudintermedius*, used as reference (CP011490.1) [54]. Finally, a Class II type BGC was predicted in *S. xyloso* C9255 which was related to the blp bacteriocin family [55].

Circular Bacteriocins

In the present study, three putative novel circular BGCs have been identified in *S. hyicus* C9581, *S. xyloso* C9255, and *S. chromogenes* C9838 isolates recovered from wild animals. The detected coding genes showed identity with the conserved domain of uberolysin, a circular bacteriocin firstly detected in *Streptococcus uberis* [56]. Comparative analysis of the BGCs of the potential circular bacteriocin carriers predicted in the present study and those used as references (*S. aureus*_WH39_CP060492, *S. uberis*_DQ650653 and *S. aureus*_pRJ101_Aureocyclicin4185_KF836421) was performed. A high identity was exhibited between one of the BGC predicted in the *S. chromogenes* C9838 and one of *S. xyloso* C9255 genomes with the operon coding for the circular bacteriocin described for *S. aureus* WH39. Additionally, these BGCs were closely related with the aureocyclicin 4185 BGC and the strongest identity was detected when compared with the BGC predicted in the *S. hyicus* C9581 genome. Finally, the other BGC predicted to code for a circular bacteriocin in *S. chromogenes* C9838 could be considered as an unrelated genetic system (Fig. S2).

Non-Ribosomal Peptides

Regarding NRPs, operons related with this type of secondary metabolites were predicted in six genomes from a wide variety of species and origins (Table 1). Genome analysis revealed high diversity between them, and no bacteriocins were detected except for the two clinical *S. lugdunensis* genomes (9%), carrying the lugdunin BGC. Interestingly, comparisons with the lugdunin coding operon revealed high identity > 95% with the first discovered lugdunin producer (*S. lugdunensis*_IVK28_CP063143.1) (Fig. S3) [57].

Thiopeptides

Previous studies by our group confirmed the production of MP1 bacteriocin by environmental staphylococcal isolates using mass-spectrometry analysis [58]. Based on this finding, we decided to further characterize these isolates by genome mining tools. Thus, the BGC for MP1 thiopeptide production were detected in sequences of the four bacteriocin-producing isolates included in this study (*S. aureus* C5802 and *S. hominis* C5835 isolates from environmental samples; two *S. sciuri* X3011 and X3041 isolates from food).

Genetic environment comparisons between the four MP1 producers included in this study and those previously described confirmed the high identity between the BGC structures of isolates from the genus *Staphylococcus* but with clear differences to the BGCs from other genera (Fig. 4). The reference sequences used for the MP1 operon analysis were as follows (isolate reference/accession number): *Bacillus safensis* (SDG14_10_1/JACEVK010000010), *Bacillus cereus* (ATCC14579/CP034551), *Listeria monocytogenes* (FDA802499/AAAXMT010000009), *Streptococcus pseudoporcinus* (NCTC5385/LR594035), *Macrococcus caseolyticus* (pBac115/KM613043), *Staphylococcus agnetis* (4244/JABULG020000014), *S. intermedius* (14S03307/MWRT01000009), *Staphylococcus felis* (F30k1271111/QKYH01000057), *S. aureus* (358/CP077876), *S. hominis* (34/CP040733), and *S. sciuri* (IMDOS72p/CAJVGN010000003). Moreover, close to 100% identity was observed among *S. intermedius* (14503307) and *S. felis* (F30k1271111), between *S. aureus* (358) and *S. hominis* (34), and between the two independent *S. sciuri* isolates. In addition, the DNA sequence analysis of the MP1 genetic environment revealed the closest relationship between the operon identified in *M. caseolyticus* (pbac115) with those found in *Staphylococcus*. The strongest differences were observed between *Staphylococcus* and *M. caseolyticus* (115) MP1 operons with those described in *Bacillus* spp. isolates (SDG14_10_1 or ATCC14579) (Fig. 4).

For deeper research of the MP1 BGC detected among the isolates included in this study, a global comparison with the NCBI database was performed. Individual BlastN analysis with each of the contigs carrying the MP1 BGC revealed a close affinity between the MP1 operon of our four carrier genomes and a complete plasmid coding for MP1 BGC of *S. hominis*_34 (CP040733.1). Moreover, the tight relationship between the entire MP1 plasmid of *S. hominis*_34 and of *S. hominis* C5835, recovered from this study, is noteworthy. In this respect, the BlastN analysis revealed a high identity and coverage percentage (> 90%), differing in 5 genes coding for ATPases or genes with unknown function. This identity was lower when comparing the MP1 BGC of the *S. aureus*




Fig. 4 Genetic environment comparison of MPI thiopeptide gene clusters of *Bacillus safensis* (SDG14_10_1/JACEVK010000010), *Bacillus cereus* (ATCC14579/CP034551), *Listeria monocytogenes* (FDA802499/AAAXMT010000009), *Streptococcus pseudoporcinus* (NCTC5385/LR594035), *Macroccoccus caseolyticus* (pBac115/KM613043), *Staphylococcus agnetis* (4244/JABULG020000014), *S. intermedius* (14S03307/

MWRT01000009), *S. felis* (F30k1271111/QKYH01000057), *S. aureus* (358/CP077876), *S. hominis* (34/CP040733), *S. sciuri* (IMDOS72p/CAJVGN010000003), *S. hominis* (C5835, this study), *S. aureus* (C5802, this study), *S. sciuri* (X3041, this study), and *S. sciuri* (X3011, this study). This figure is in color in the online version but in black and white in the print version




Fig. 5 Differences at the protein level of the structural gene coding for the thiopeptide MP1: *B. safensis* (SDG14_10_1), *B. cereus* (ATCC14579), *L. monocytogenes* (FDA802499), *S. pseudoporcinus* (NCTC5385), *M. caseolyticus* (pBac115), *S. agnetis* (4244), *S. intermedius* (14S03307), *S. felis* (F30k1271111), *S. aureus* (358), *S.*

hominis (34), *S. sciuri* (IMDOS72p), *S. hominis* (C5835, this study), *S. aureus* (C5802, this study), *S. sciuri* (X3041, this study), *S. sciuri* (X3011, this study). This figure is in color in the online version but in black and white in the print version

C5802 isolate with *S. hominis*_34. However, high identity was observed between the *S. aureus* C5802 and *S. hominis* C5835 MP1 operons, also when comparing the flanking areas so it can be suggested that these two isolates could share a common genetic element. Finally, the presence of identical additional genes upstream of the MP1 operon in the two *S. sciuri* from this study and the selected reference (MP1 coding plasmid of *S. sciuri* IMDOS72) should be mentioned. Further analysis should be carried out to confirm the relation of the four MP1 BGC detected in this study to mobile genetic elements such as plasmids and their transmission capability.

Moreover, differences in the MP1 structural gene were studied at the protein level (Fig. 5). Regarding the amino acid sequence alignment, it is important to note that *Staphylococcus* isolates were all identical when considering the MP1 structural gene, except for *S. agnetis* from the database which presented two amino acid changes in the leader peptide. Moreover, the high identity between *Staphylococcus* and *M. caseolyticus* sequences was outstanding, which only differs in a few amino acids within the leader peptide. On the other hand, the comparison with the other isolates included in this study revealed huge differences in the leader peptide sequences and even punctual amino acid exchanges and insertions within the mature peptides of *L. monocytogenes* and *S. pseudoporcinus*, likely affecting the structure and/or antimicrobial activity of their bacteriocins.

Discussion

The new genome mining tools offer an important technological resource in the discovery of novel natural products based either on the detection of bacteriocin structural genes or other bacteriocin-associated genes [59]. In this study, ~82% of the investigated bacteriocin-producing *Staphylococcus* isolates (18 out of 22 genomes) carried at least one bacteriocin coding gene among which a large diversity was observed independently of isolate origin (human, pet, wild animals, food, and environmental). Here, five types of BGCs were identified among 18 bacteriocin-producing isolates, coding for bacteriocins of class I (lanthipeptides), class II (lactococcin972, bacSp222, and blp family bacteriocin), class IV (circular bacteriocins), NRP, and thiopeptides. Interesting relationship between related *Staphylococcus* species and the type of bacteriocin has been observed.

Focusing on lanthipeptides, they are characterized by their small (< 5 kDa) size, their post-translational modifications and they contain lanthionine or β -methylanthionine residues in their structure [60, 61]. In the present work, seven lanthipeptide-like BGCs have been described among the 22 studied genomes, and genomic comparisons with those bacteriocin clusters previously referred in the literature allowed us to classify these putative new bacteriocins into class I. Upon lanthipeptide BGC identification

among bacteriocin-producing staphylococci, bioinformatic analyses indicated the highest identity of the bacteriocins predicted in this study to the reference ones: (i) the BSA bacteriocin discovered from an MRSA strain involved in community-acquired infections [62]; (ii) the bacCH91 bacteriocin reported by Wladyka and colleagues in 2013 [63], although they only reported the structural gene; (iii) the epilancin15X antimicrobial peptide firstly detected in a clinical *S. epidermidis* isolate [64]. These results showed a close relationship between the putative structural genes detected in the newly investigated isolates and those previously reported, which suggests the detection of new variants of both BSA and epilancin15X bacteriocins. In addition, a lanthipeptide-like BGC type (V) was detected in the *S. hyicus* C9585 genome showing identity with those recently reported elsewhere [65, 66].

Interestingly, the BGCs detected in *S. warneri* and *S. epidermidis* bacteriocin-producing isolates seemed to be putative new lanthipeptide-like bacteriocins. Lanthipeptide BGC are characterized by the presence of a structural gene which is post-translationally modified by dehydratase, phosphatase, glycosidase, cyclase, oxidase enzymes, and ABC transporters [18]. Moreover, the presence of a C-terminal core region in the structural gene of *S. simulans* C9832 suggests that this operon could code for a type I lanthipeptide, while the one detected in *S. epidermidis* X3009 revealed higher identity to pep5 bacteriocin (Figs. 2 and 3). Future work will focus on the optimization of the extraction conditions to obtain inhibition in the well diffusion assay and to finally confirm the proteinaceous nature of the inhibitory compound and to decipher the peptide structure. Moreover, the use of heterologous expression systems will be important for future validation of the in silico screening studies [67–69].

Focusing on class II bacteriocins, lactococcin-like clusters were identified in six out of the 22 bacteriocin-producing isolates included. These bacteriocins are described as linear non-pediocin-like molecules. Among them, lactococcin972 produced by *Lactococcus lactis* subsp. *lactis* IPLA 972 is the most representative and frequent [70]. Figure S1 shows a lactococcin-like operon comparison, and all clusters contained the precursor of lactococcin972 domain.

In addition, the bacSp222 bacteriocin of class II has also been detected in this study. It was the first reported bacteriocin produced by *S. pseudintermedius* [54], and its genetic cluster organization revealed identity to aureocin A53. Moreover, bacsp222 is characterized by its bactericidal activity and virulence capacity that acts in modulating the immune system of the host [54].

The recently proposed class IV bacteriocins are circular peptides, formed by the post-translational covalent linkage between their carboxy and amino termini [71]. Currently, aureocyclicin 4185 from *S. aureus* 4185 is the first circular bacteriocin of *Staphylococcus* included in this category [72]. In this respect,

we identified here two putative novel BGCs encoding circular bacteriocins of the circularin A/uberolysin family without identity to the previously reported circular staphylococci (Fig. S2). Moreover, the *S. hyicus* C9581 isolate seems to carry the aureocyclicin 4185 bacteriocin coding genes.

Finally, cyclic lugdunin, which was the first NRP described among staphylococci and is considered as an independent class of antibacterials, was detected in two clinical *S. lugdunensis* isolates of the present work. Likewise, Zipperer and collaborators discovered lugdunin in a nasal *S. lugdunensis* isolate with antimicrobial activity against a wide range of Gram-positive bacteria including methicillin-resistant *S. aureus* (MRSA) [57]. The NRPS operon is about 30 kbp and encodes four genes (*lugABCD*) for the biosynthesis of lugdunin. Not all *S. lugdunensis* produce antimicrobial peptides although the lugdunin BGC seems to be conserved in this species and interestingly, the GC-content indicates horizontal transfer to *S. lugdunensis* from another bacterial species [57].

Regarding the bacteriocin classification proposed by de Freire Bastos in 2020, thiopeptides could be considered as a new class of staphylococci (Class VI) [13]. Thiopeptides are sulfur-containing, ribosomally produced and highly post-translationally modified peptides with strong inhibitory and competitive potential [73, 74]. Most likely, MP1 (originally designated micrococin) was firstly discovered in a strain of *Micrococcus* by its activity against *Mycobacterium tuberculosis* and it was characterized as a hydrophobic and heat-stable molecule with high activity against a wide range of Gram-positive bacteria [75]. So far, MP1 has been isolated from different genera, including *Micrococcus*, *Staphylococcus*, *Streptococcus*, and *Bacillus* spp., and origins (food, humans, and animals).

Focusing on *Staphylococcus*, MP1 production has been reported for isolates of *S. equorum* from cheese [76], *S. epidermidis* [77], *S. felis* from cats [78], *S. hominis* recovered from human skin [79], *S. sciuri* [29], and recently, *S. aureus* [30]. In the present study, MP1 production has been verified at the genetic level among four bacteriocin-producing isolates (~18%) with high antimicrobial activity against both MSSA and MRSA: two *S. aureus* and *S. hominis* isolates from environmental samples (river water) and two *S. sciuri* isolates recovered from raw meat chicken.

Comparison of the MP1 BGCs (Fig. 4) revealed major genetic differences between the staphylococcal isolates and *Bacillus*. First, the number of structural genes can differ between both genera (one for *Staphylococcus* and up to four for *Bacillus cereus*). Next, the staphylococcal strains appear to produce only one product [77], while a mixture of similar thiopeptides with different post-translational modifications [thiocillin I, II, III, MP1 and micrococin P2 (MP2)] have been reported among *Bacillus* [80]. Only after horizontal gene transfer of the MP1 BGC into *S. aureus* RN4220, a yet uncharacterized by-product of MP1 could be detected [30]. In this study,

the strong capacity of MP1 to force RN4220 to change its metabolic capacity via *citZ* mutation was highlighted. In addition, the comparison of the MP1 BGCs included in this study illustrates the high identity between the *Staphylococcus* isolates, especially when considering the same species and with *M. caseolyticus*.

Most staphylococcal BGCs appear to be associated with mobile genetic elements such as plasmids, transposons, IS-elements, or chromosomal islands [28, 30, 57]. Hence, BGCs can be transferred between strains and lineages and are important genetic determinants of competitive fitness within a given habitat [30]. Due to the great diversity of staphylococcal isolates and origins detected among our bacteriocin-producing isolates and the high frequency of MP1 carrier detection, the occurrence of certain mechanisms of BGC transfer could be assumed as mentioned above.

The analysis of the genomes included in this study allowed us to identify a wide diversity of BGCs and more concretely, the comparison of the genetic environment of MP1 revealed identity to the reference plasmids of *S. hominis* and *S. sciuri*, indicating that the BGCs are plasmid-encoded. However, although the detection of *rep* sequences by PlasmidFinder, third-generation sequencing technologies, such as PacBio [81] or Oxford Nanopore (ONT) [82] instruments are recommended to confirm the presence of transferrable BGC.

Conclusion

In conclusion, our findings revealed a great abundance and diversity of bacteriocin gene clusters including unique systems and unfrequently detected among staphylococcal genomes. In this respect, the genus *Staphylococcus* and specially CoNS isolates have been confirmed as a valuable source of new peptide structures with promising functionalities for treatment and prevention. Moreover, the OneHealth perspective should be accentuated as a good perspective for further research on the alternatives for the AMR crisis.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s12602-023-10119-w>.

Acknowledgements The authors acknowledge support by the High Performance and Cloud Computing Group at the Zentrum für Datenverarbeitung of the University of Tübingen, the state of Baden-Württemberg through bwHPC, and the German Research Foundation (DFG) through grant no INST 37/935-1 FUGG.

Author Contribution C.T., B.K., A.P., R.F.-F., and C.L. contributed to the design of the study. C.T., B.K., A.P., and C.L. contributed to the general supervision of the study. A.M.A.E. and M.T. carried out the processing of the sequences. R.F.-F., A.M.A.E., and A.M. collaborated in the data analysis and R. F.-F. developed the first version of the manuscript. C.T., C.L., and B.K. made the first revision of the manuscript. C.T., M.Z. and A.P. contributed to project funding. All authors revised

the different versions of the manuscript, read, and agreed to the submitted version of the manuscript.

Funding Open Access funding provided thanks to the CRUE-CSIC agreement with Springer Nature. This study was funded by the project PID2019-106158RB-I00 from the Ministry of Spain (MCIN/AEI/10.13039/501100011033). A.P. acknowledges grants from Deutsche Forschungsgemeinschaft (DFG) TRR156 (project ID 246807620); TRR261 (project ID 398967434); and Cluster of Excellence EXC2124 Controlling Microbes to Fight Infection (CMFI) (project ID 390838134). R.F.-F. has a predoctoral contract FPU from the Ministry of Science, Innovation and Universities of Spain (FPU18/05438) and a training fellowship for predoctoral student exchanges in international research centers (Ministry of Spain, EST21/00631).

Data Availability All data generated or analyzed during this study are available within this paper and its supplementary information files. References [75–79] corresponds to Supplementary material (Table S2). Illumina reads for 21 staphylococcal isolates included in this study can be found at the BioProject PRJNA974190 and C5802 is registered with ERS659514 accession number (ENA). The respective BioSample numbers are indicated in Supplementary Table S2.

Declarations

Ethics Approval This article does not contain any studies with human participants or animals performed by any of the authors.

Competing Interests The authors declare no competing interests.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Murray CJL, Ikuta KS, Sharara F, Swetschinski L, Robles Aguilar G, Gray A, Han C, Bisignano C, Rao P, Wool E, Johnson SC, Browne AJ, Chipeta MG, Fell F, Hackett S, Haines-Woodhouse G, Kashef Hamadani BH, Kumaran EAP, McManigal B, Achalapong S, Agarwal R, Akech S, Albertson S, Amuasi J, Andrews J, Aravkin A, Ashley E, Babin F-X, Bailey F, Baker S, Basnyat B, Bekker A, Bender R, Berkley JA, Bethou A, Bielicki J, Boonkasidecha S, Bukosia J, Carvalho C, Castañeda-Orjuela C, Chansamouth V, Chaurasia S, Churchiù S, Chowdhury F, Clotaire Donatien R, Cook AJ, Cooper B, Cressey TR, Criollo-Mora E, Cunningham M, Darboe S, Day NPJ, De Luca M, Dokova K, Dramowski A, Dunachie SJ, Duong Bich T, Eckmanns T, Eibach D, Emami A, Feasey N, Fisher-Pearson N, Forrest K, Garcia C, Garrett D, Gastmeier P, Giref AZ, Greer RC, Gupta V, Haller S, Haselbeck A, Hay SI, Holm M, Hopkins S, Hsia Y, Iregbu KC, Jacobs J, Jarovsky D, Javanmardi F, Jenney AWJ, Khorana M, Khusuwan S, Kissoon N, Kobeissi E, Kostyanov T, Krapp F, Krumkamp R, Kumar A, Kyu HH, Lim C, Lim K, Limmathurotsakul D, Loftus MJ, Lunn M, Ma J, Manoharan A,

- Marks F, May J, Mayxay M, Mturi N, Munera-Huertas T, Musicha P, Musila LA, Mussi-Pinhata MM, Naidu RN, Nakamura T, Nanavati R, Nangia S, Newton P, Ngoun C, Novotney A, Nwakanma D, Obiero CW, Ochoa TJ, Olivas-Martinez A, Olliaro P, Ooko E, Ortiz-Brizuela E, Ounchanum P, Pak GD, Paredes JL, Peleg AY, Perrone C, Phe T, Phommasone K, Plakkal N, Ponce-de-Leon A, Raad M, Ramdin T, Rattanavong S, Riddell A, Roberts T, Robotham JV, Roca A, Rosenthal VD, Rudd KE, Russell N, Sader HS, Saengchan W, Schnell J, Scott JAG, Seekaew S, Sharland M, Shivamallappa M, Sifuentes-Osornio J, Simpson AJ, Steenkeste N, Stewardson AJ, Stoeva T, Tasak N, Thaiprakong A, Thwaites G, Tigoi C, Turner C, Turner P, Van Doorn HR, Velaphi S, Vongpradith A, Vongsouvath M, Vu H, Walsh T, Watson JL, Waner S, Wangrangsimakul T, Wannapinij P, Wozniak T, Young Sharma TEMW, Yu KC, Zheng P, Sartorius B, Lopez AD, Stergachis A, Moore C, Dolecek C, Naghavi M (2022) Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. *Lancet* 399:629–655. [https://doi.org/10.1016/S0140-6736\(21\)02724-0](https://doi.org/10.1016/S0140-6736(21)02724-0)
2. Ovchinnikov KV, Kranjec C, Telke A, Kjos M, Thorstensen T, Scherer S, Carlsen H, Diep DB (2021) A strong synergy between the thiopeptide bacteriocin micrococcin P1 and rifampicin against MRSA in a murine skin infection model. *Front Immunol* 12:676534. <https://doi.org/10.3389/fimmu.2021.676534>
 3. Aljeldah MM (2022) Antimicrobial resistance and its spread is a global threat. *Antibiotics* 11:1082. <https://doi.org/10.3390/antibiotics11081082>
 4. Calixto JB (2019) The role of natural products in modern drug discovery. *An Acad Bras Ciênc* 91:e20190105. <https://doi.org/10.1590/0001-3765201920190105>
 5. Cragg GM, Newman DJ, Snader KM (1997) Natural products in drug discovery and development. *J Nat Prod* 60:52–60. <https://doi.org/10.1021/np9604893>
 6. Strohl WR (2000) The role of natural products in a modern drug discovery program. *Drug Discovery Today* 5:39–41. [https://doi.org/10.1016/S1359-6446\(99\)01443-9](https://doi.org/10.1016/S1359-6446(99)01443-9)
 7. Morton JT, Freed SD, Lee SW, Friedberg I (2015) A large scale prediction of bacteriocin gene blocks suggests a wide functional spectrum for bacteriocins. *BMC Bioinformatics* 16:381. <https://doi.org/10.1186/s12859-015-0792-9>
 8. Arnison PG, Bibb MJ, Bierbaum G, Bowers AA, Bugni TS, Bulaj G, Camarero JA, Campopiano DJ, Challis GL, Clardy J, Cotter PD, Craik DJ, Dawson M, Dittmann E, Donadio S, Dorrestein PC, Entian K-D, Fischbach MA, Garavelli JS, Göransson U, Gruber CW, Haft DH, Hemscheidt TK, Hertweck C, Hill C, Horswill AR, Jaspars M, Kelly WL, Klinman JP, Kuipers OP, Link AJ, Liu W, Marahiel MA, Mitchell DA, Moll GN, Moore BS, Müller R, Nair SK, Nes IF, Norris GE, Olivera BM, Onaka H, Patchett ML, Piel J, Reaney MJT, Rebuffat S, Ross RP, Sahl HG, Schmidt EW, Selsted ME, Severinov K, Shen B, Sivonen K, Smith L, Stein T, Süßmuth RD, Tagg JR, Tang GL, Truman AW, Vederas JC, Walsh CT, Walton JD, Wenzel SC, Willey JM, Van Der Donk WA (2013) Ribosomally synthesized and post-translationally modified peptide natural products: overview and recommendations for a universal nomenclature. *Nat Prod Rep* 30:108–160. <https://doi.org/10.1039/C2NP20085F>
 9. Azevedo AC, Bento CBP, Ruiz JC, Queiroz MV, Mantovani HC (2015) Distribution and genetic diversity of bacteriocin gene clusters in rumen microbial genomes. *Appl Environ Microbiol* 81:7290–7304. <https://doi.org/10.1128/AEM.01223-15>
 10. Carson DA, Barkema HW, Naushad S, De Buck J (2017) Bacteriocins of non-aureus staphylococci isolated from bovine milk. *Appl Environ Microbiol* 83:e01015–e1017. <https://doi.org/10.1128/AEM.01015-17>
 11. Yap PG, Lai ZW, Tan JS (2022) Bacteriocins from lactic acid bacteria: purification strategies and applications in food and medical industries: a review. *Beni-Suef Univ J Basic Appl Sci* 11:51. <https://doi.org/10.1186/s43088-022-00227-x>
 12. Bennalack PR, Bewley KD, Burlingame MA, Robison RA, Miller SM, Griffiths JS (2016) Reconstitution and minimization of a micrococcin biosynthetic pathway in *Bacillus subtilis*. *J Bacteriol* 198:2431–2438. <https://doi.org/10.1128/JB.00396-16>
 13. Silva CCG, Silva SPM, Ribeiro SC (2018) Application of bacteriocins and protective cultures in dairy food preservation. *Front Microbiol* 9:594. <https://doi.org/10.3389/fmicb.2018.00594>
 14. Alvarez-Sieiro P, Montalbán-López M, Mu D, Kuipers OP (2016) Bacteriocins of lactic acid bacteria: extending the family. *Appl Microbiol Biotechnol* 100:2939–2951. <https://doi.org/10.1007/s00253-016-7343-9>
 15. de Freire Bastos MDC, Miceli De Farias F, Carlin Fagundes P, Varella Coelho ML (2020) Staphylococci: an update on antimicrobial peptides produced by staphylococci and their diverse potential applications. *Appl Microbiol Biotechnol* 104:10339–10368. <https://doi.org/10.1007/s00253-020-10946-9>
 16. Newstead LL, Varjonen K, Nuttall T, Paterson GK (2020) Staphylococcal-produced bacteriocins and antimicrobial peptides: their potential as alternative treatments for staphylococcus aureus infections. *Antibiotics* 9:40. <https://doi.org/10.3390/antibiotics9020040>
 17. Li JWH, Vederas JC (2009) Drug discovery and natural products: end of an era or an endless frontier? *Science* 325:161–165. <https://doi.org/10.1126/science.1168243>
 18. Torres Salazar BO, Heilbronner S, Peschel A, Krismer B (2021) Secondary metabolites governing microbiome interaction of staphylococcal pathogens and commensals. *Microb Physiol* 31:198–216. <https://doi.org/10.1159/000517082>
 19. Heilbronner S, Krismer B, Brötz-Oesterhelt H, Peschel A (2021) The microbiome-shaping roles of bacteriocins. *Nat Rev Microbiol* 19:726–739. <https://doi.org/10.1038/s41579-021-00569-w>
 20. Gradisteanu Pircalabioru G, Popa LI, Marutescu L, Gheorghe I, Popa M, Czobor Barbu I, Cristescu R, Chifiriuc MC (2021) Bacteriocins in the era of antibiotic resistance: rising to the challenge. *Pharmaceutics* 13:196. <https://doi.org/10.3390/pharmaceutics13020196>
 21. Mercado V, Olmos J (2022) Bacteriocin production by bacillus species: isolation, characterization, and application. *Probiotics Antimicro Prot* 14:1151–1169. <https://doi.org/10.1007/s12602-022-09966-w>
 22. Gontijo MTP, Ramia NE, Dijamentiuk A, Elfassy A, Taha S, Mangavel C, Revol-Junelles AM, Borges F (2022) Mining biosynthetic gene clusters in *Carnobacterium maltaromaticum* by interference competition network and genome analysis. *Microorganisms* 10:1794. <https://doi.org/10.3390/microorganisms10091794>
 23. Ongey EL, Neubauer P (2016) Lanthipeptides: chemical synthesis versus in vivo biosynthesis as tools for pharmaceutical production. *Microb Cell Fact* 15:97. <https://doi.org/10.1186/s12934-016-0502-y>
 24. Simons A, Alhanout K, Duval RE (2020) Bacteriocins, antimicrobial peptides from bacterial origin: overview of their biology and their impact against multidrug-resistant bacteria. *Microorganisms* 8:639. <https://doi.org/10.3390/microorganisms8050639>
 25. Bagley MC, Dale JW, Merritt EA, Xiong X (2005) Thiopeptide antibiotics. *Chem Rev* 105:685–714. <https://doi.org/10.1021/cr0300441>
 26. Fernandes A, Jobby R (2022) Bacteriocins from lactic acid bacteria and their potential clinical applications. *Appl Biochem Biotechnol* 194:4377–4399. <https://doi.org/10.1007/s12010-022-03870-3>
 27. Bosák J, Hrala M, Micenkova L, Šmajš D (2021) Non-antibiotic antibacterial peptides and proteins of *Escherichia coli*: efficacy and potency of bacteriocins. *Expert Rev Anti Infect Ther* 19:309–322. <https://doi.org/10.1080/14787210.2020.1816824>
 28. Janek D, Zipperer A, Kulik A, Krismer B, Peschel A (2016) High frequency and diversity of antimicrobial activities produced by nasal *Staphylococcus* strains against bacterial competitors. *PLoS Pathog* 12:e1005812. <https://doi.org/10.1371/journal.ppat.1005812>
 29. Van Der Veken D, Hollanders C, Verce M, Michiels C, Ballet S, Weckx S, Leroy F (2022) Genome-based characterization of

- a plasmid-associated micrococcin P1 biosynthetic gene cluster and virulence factors in *Mammaliicoccus sciuri* IMDO-S72. *Appl Environ Microbiol* 88:e02088–e2121. <https://doi.org/10.1128/aem.02088-21>
30. Krauss S, Harbig TA, Rapp J, Schaeffle T, Franz-Wachtel M, Reetz L, Elsherbini AMA, Macek B, Grond S, Link H, Nieselt K, Krismer B, Peschel A, Heilbronner S (2023) Horizontal transfer of bacteriocin biosynthesis genes requires metabolic adaptation to improve compound production and cellular fitness. *Microbiol Spectr* 11:e03176–e3222. <https://doi.org/10.1128/spectrum.03176-22>
 31. Fernández-Fernández R, Lozano C, Eguizábal P, Ruiz-Ripa L, Martínez-Álvarez S, Abdullahi IN, Zarazaga M, Torres C (2022) Bacteriocin-like inhibitory substances in staphylococci of different origins and species with activity against relevant pathogens. *Front Microbiol* 13:870510. <https://doi.org/10.3389/fmicb.2022.870510>
 32. Fernández-Fernández R, Lozano C, Ruiz-Ripa L, Robredo B, Azcona-Gutiérrez JM, Alonso CA, Aspiroz C, Zarazaga M, Torres C (2022) Antimicrobial resistance and antimicrobial activity of *Staphylococcus lugdunensis* obtained from two Spanish hospitals. *Microorganisms* 10:1480. <https://doi.org/10.3390/microorganisms10081480>
 33. Fernández-Fernández R, Abdullahi IN, González-Azcona C, Ulloa A, Martínez A, García-Vela S, Höfle U, Zarazaga M, Lozano C, Torres C (2023) Detection of antimicrobial producing *Staphylococcus* from migratory birds: potential role in nasotracheal microbiota modulation. *Front Microbiol* 14:1144975. <https://doi.org/10.3389/fmicb.2023.1144975>
 34. Chun J, Oren A, Ventosa A, Christensen H, Arahal DR, Da Costa MS, Rooney AP, Yi H, Xu XW, De Meyer S, Trujillo ME (2018) Proposed minimal standards for the use of genome data for the taxonomy of prokaryotes. *Int J Syst Evol Microbiol* 68:461–466. <https://doi.org/10.1099/ijsem.0.002516>
 35. Madhaiyan M, Wirth JS, Saravanan VS (2020) Phylogenomic analyses of the *Staphylococcaceae* family suggest the reclassification of five species within the genus *Staphylococcus* as heterotypic synonyms, the promotion of five subspecies to novel species, the taxonomic reassignment of five *Staphylococcus* species to *Mammaliicoccus* gen. nov, and the formal assignment of *Nosocomiicoccus* to the family *Staphylococcaceae*. *Int J Syst Evol Microbiol* 70:5926–5936. <https://doi.org/10.1099/ijsem.0.004498>
 36. Andrews S (2010) FastQC: A quality control tool for high throughput sequence data. Available online at: <http://www.bioinformatics.babraham.ac.uk/projects/fastqc>
 37. Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>
 38. Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A (2020) Using SPAdes de novo assembler. *Curr Protoc Bioinform* 70. <https://doi.org/10.1002/cpbi.102>
 39. Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q, Wortman J, Young SK, Earl AM (2014) Pilon: an integrated tool for comprehensive microbial variant detection and genome assembly improvement. *PLoS ONE* 9:e112963. <https://doi.org/10.1371/journal.pone.0112963>
 40. Seemann T (2019). Shovill pipeline. Available online at: <https://github.com/tseemann/shovill>
 41. Seemann T (2014) Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>
 42. Stamatakis A (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30:1312–1313. <https://doi.org/10.1093/bioinformatics/btu033>
 43. Asnicar F, Thomas AM, Beghini F, Mengoni C, Manara S, Manghi P, Zhu Q, Bolzan M, Cumbo F, May U, Sanders JG, Zolfo M, Kopylova E, Pasolli E, Knight R, Mirarab S, Huttenhower C, Segata N (2020) Precise phylogenetic analysis of microbial isolates and genomes from metagenomes using PhyloPhlAn 3.0. *Nat Commun* 11:2500. <https://doi.org/10.1038/s41467-020-16366-7>
 44. Xu S, Li L, Luo X, Chen M, Tang W, Zhan L, Dai Z, Lam TT, Guan Y, Yu G (2022) *Ggtree*: a serialized data object for visualization of a phylogenetic tree and annotation data. *iMeta* 1. <https://doi.org/10.1002/imt2.56>
 45. R Core Team (2018) R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria
 46. Technical University of Denmark (2023) Center of genomic epidemiology server. Available online at: <http://www.genomicepidemiology.org>
 47. Blin K, Shaw S, Augustijn HE, Reitz ZL, Biermann F, Alanjary M, Fetter A, Terlouw BR, Metcalf WW, Helfrich EJN, van Wezel GP, Medema MH, Weber T (2023) antiSMASH 7.0: new and improved predictions for detection, regulation, chemical structures and visualisation. *Nucleic Acids Res* gkad344. <https://doi.org/10.1093/nar/gkad344>
 48. van Heel A, de Jon A, Song C, Viel JH, Kok J, Kuipers OP (2018) BAGEL4: a user-friendly web server to thoroughly mine RiPPs and bacteriocins. *Nucleic Acids Res* 46:W278–W281. <https://doi.org/10.1093/nar/gky383>
 49. Gilchrist CLM, Booth TJ, Van Wersch B, Van Grieken L, Medema MH, Chooi YH (2021) cblaster: a remote search tool for rapid identification and visualization of homologous gene clusters. *Bioinformatics Adv* 1:vbab016. <https://doi.org/10.1093/bioadv/vbab016>
 50. Gilchrist CLM, Chooi YH (2021) clinker & clustermap.js: automatic generation of gene cluster comparison figures. *Bioinformatics* 37:2473–2475. <https://doi.org/10.1093/bioinformatics/btab007>
 51. Waterhouse AM, Procter JB, Martin DMA, Clamp M, Barton GJ (2009) Jalview Version 2—a multiple sequence alignment editor and analysis workbench. *Bioinformatics* 25:1189–1191. <https://doi.org/10.1093/bioinformatics/btp033>
 52. Madeira F, Pearce M, Tivey ARN, Basutkar P, Lee J, Edbali O, Madhusoodanan N, Kolesnikov A, Lopez R (2022) Search and sequence analysis tools services from EMBL-EBI in 2022. *Nucleic Acids Res* 50:W276–W279. <https://doi.org/10.1093/nar/gkac240>
 53. Kim PI, Sohng JK, Sung C, Joo HS, Kim EM, Yamaguchi T, Park D, Kim BG (2010) Characterization and structure identification of an antimicrobial peptide, hominicin, produced by *Staphylococcus hominis* MBBL 2–9. *Biochem Biophys Res Commun* 399:133–138. <https://doi.org/10.1016/j.bbrc.2010.07.024>
 54. Władyska B, Piejko M, Bzowska M, Pieta P, Krzysiek M, Mazurek Ł, Guevara-Lora I, Bukowski M, Sabat AJ, Friedrich AW, Bonar E, Międzobrodzki J, Dubin A, Mak P (2015) A peptide factor secreted by *Staphylococcus pseudintermedius* exhibits properties of both bacteriocins and virulence factors. *Sci Rep* 5:14569. <https://doi.org/10.1038/srep14569>
 55. Fontaine L, Boutry C, Guédon E, Guillot A, Ibrahim M, Grossiord B, Hols P (2007) Quorum-sensing regulation of the production of Blp bacteriocins in *Streptococcus thermophilus*. *J Bacteriol* 189:7195–7205. <https://doi.org/10.1128/JB.00966-07>
 56. Lasagno M, Navarro MDLA, Moliva M, Reinoso E (2019) Screening of bacteriocin associated genes of *Streptococcus uberis* strains. *Heliyon* 5:e02393. <https://doi.org/10.1016/j.heliyon.2019.e02393>
 57. Zipperer A, Konnerth MC, Laux C, Berscheid A, Janek D, Weidenmaier C, Burian M, Schilling NA, Slavetinsky C, Marschal M, Willmann M, Kalbacher H, Schittek B, Brötz-Oesterhelt H, Grond S, Peschel A, Krismer B (2016) Human commensals producing a novel antibiotic impair pathogen colonization. *Nature* 535:511–516. <https://doi.org/10.1038/nature18634>

58. Fernández-Fernández R, Lozano C, Fernández-Pérez R, Zarazaga M, Peschel A, Krismser B, Torres C (2023, under revision) Detection of micrococcin MP1 in commensal and environmental staphylococcal isolates with high antimicrobial activity against MRSA. *Int J Antimicrob Agents* (in revision)
59. Egan K, Field D, Ross RP, Cotter PD, Hill C (2018) In silico prediction and exploration of potential bacteriocin gene clusters within the bacterial genus *Geobacillus*. *Front Microbiol* 9:2116. <https://doi.org/10.3389/fmicb.2018.02116>
60. Jack RW, Tagg JR, Ray B (1995) Bacteriocins of gram-positive bacteria. *Microbiol Rev* 59:171–200. <https://doi.org/10.1128/mr.59.2.171-200.1995>
61. McAuliffe O, Ross RP, Hill C (2001) Lantibiotics: structure, biosynthesis and mode of action. *FEMS Microbiol Rev* 25:285–308. <https://doi.org/10.1111/j.1574-6976.2001.tb00579.x>
62. Daly KM, Upton M, Sandiford SK, Draper LA, Wescombe PA, Jack RW, O'Connor PM, Rossney A, Götz F, Hill C, Cotter PD, Ross RP, Tagg JR (2010) Production of the Bsa lantibiotic by community-acquired *Staphylococcus aureus* strains. *J Bacteriol* 192:1131–1142. <https://doi.org/10.1128/JB.01375-09>
63. Wladyka B, Wielebska K, Wloka M, Bochenska O, Dubin G, Dubin A, Mak P (2013) Isolation, biochemical characterization, and cloning of a bacteriocin from the poultry-associated *Staphylococcus aureus* strain CH-91. *Appl Microbiol Biotechnol* 97:7229–7239. <https://doi.org/10.1007/s00253-012-4578-y>
64. Ekkelenkamp MB, Hanssen M, Danny Hsu S-T, De Jong A, Milatovic D, Verhoef J, Van Nuland NAJ (2005) Isolation and structural characterization of epilancin 15X, a novel lantibiotic from a clinical strain of *Staphylococcus epidermidis*. *FEBS Lett* 579:1917–1922. <https://doi.org/10.1016/j.febslet.2005.01.083>
65. Pei Z-F, Zhu L, Sarkisian R, Van Der Donk WA, Nair SK (2022) Class V Lanthipeptide cyclase directs the biosynthesis of a stapled peptide natural product. *J Am Chem Soc* 144:17549–17557. <https://doi.org/10.1021/jacs.2c06808>
66. Xu M, Zhang F, Cheng Z, Bashiri G, Wang J, Hong J, Wang Y, Xu L, Chen X, Huang S, Lin S, Deng Z, Tao M (2020) Functional genome mining reveals a class V lanthipeptide containing a D-amino acid introduced by an F₄₂₀H₂-dependent reductase. *Angew Chem Int Ed* 59:18029–18035. <https://doi.org/10.1002/anie.202008035>
67. Piper C, Hill C, Cotter PD, Ross RP (2011) Bioengineering of a nisin A-producing *Lactococcus lactis* to create isogenic strains producing the natural variants nisin F, Q and Z: Comparing natural Nisin variants. *Microb Biotechnol* 4:375–382. <https://doi.org/10.1111/j.1751-7915.2010.00207.x>
68. Van Heel AJ, Kloosterman TG, Montalban-Lopez M, Deng J, Plat A, Baudu B, Hendriks D, Moll GN, Kuipers OP (2016) Discovery, production and modification of five novel lantibiotics using the promiscuous nisin modification machinery. *ACS Synth Biol* 5:1146–1154. <https://doi.org/10.1021/acssynbio.6b00033>
69. Mesa-Pereira B, O'Connor PM, Rea MC, Cotter PD, Hill C, Ross RP (2017) Controlled functional expression of the bacteriocins pediocin PA-1 and bactofoenicin A in *Escherichia coli*. *Sci Rep* 7:3069. <https://doi.org/10.1038/s41598-017-02868-w>
70. Martínez B, Fernández M, Suárez JE, Rodríguez A (1999) Synthesis of lactococcin 972, a bacteriocin produced by *Lactococcus lactis* IPLA 972, depends on the expression of a plasmid-encoded bicistronic operon The GenBank accession number for the sequence reported in this paper is AJ002203. *Microbiology* 145:3155–3161. <https://doi.org/10.1099/00221287-145-11-3155>
71. Van Belkum MJ, Martin-Visscher LA, Vederas JC (2011) Structure and genetics of circular bacteriocins. *Trends Microbiol* 19:411–418. <https://doi.org/10.1016/j.tim.2011.04.004>
72. Potter A, Ceotto H, Coelho MLV, Guimarães AJ, Bastos MDCDF (2014) The gene cluster of aureocyclin 4185: the first cyclic bacteriocin of *Staphylococcus aureus*. *Microbiology* 160:917–928. <https://doi.org/10.1099/mic.0.075689-0>
73. Eijssink VGH, Axelsson L, Diep DB, Håvarstein LS, Holo H, Nes IF (2002) Production of class II bacteriocins by lactic acid bacteria; an example of biological warfare and communication. *Antonie Van Leeuwenhoek* 81:639–654. <https://doi.org/10.1023/a:1020582211262>
74. Acedo JZ, Chiorean S, Vederas JC, Van Belkum MJ (2018) The expanding structural variety among bacteriocins from Gram-positive bacteria. *FEMS Microbiol Rev* 42:805–828. <https://doi.org/10.1093/femsre/fuy033>
75. Su TL (1948) Micrococcin, an antibacterial substance formed by a strain of *Micrococcus*. *Br J Exp Pathol* 29:473–481
76. Carnio MC, Höltzel A, Rudolf M, Henle T, Jung G, Scherer S (2000) The macrocyclic peptide antibiotic micrococcin P₁ is secreted by the food-borne bacterium *Staphylococcus equorum* WS 2733 and inhibits *Listeria monocytogenes* on soft cheese. *Appl Environ Microbiol* 66:2378–2384. <https://doi.org/10.1128/AEM.66.6.2378-2384.2000>
77. Bennallack PR, Burt SR, Heder MJ, Robison RA, Griffiths JS (2014) Characterization of a novel plasmid-borne thiopeptide gene cluster in *Staphylococcus epidermidis* strain 115. *J Bacteriol* 196:4344–4350. <https://doi.org/10.1128/JB.02243-14>
78. O'Neill AM, Worthing KA, Kulkarni N, Li F, Nakatsuji T, McGrosso D, Mills RH, Kalla G, Cheng JY, Norris JM, Pogliano K, Pogliano J, Gonzalez DJ, Gallo RL (2021) Antimicrobials from a feline commensal bacterium inhibit skin infection by drug-resistant *S. pseud-intermedius*. *eLife* 10:e66793. <https://doi.org/10.7554/eLife.66793>
79. Yao L, Liu Yuanzhen DuZ, Zhang L, Chen J, Shen Z, Liu Q, Qin J, Lv H, Wang H, He L, Liu J, Huang Q, Sun Y, Otto M, Li M (2020) Skin microbiota analysis-inspired development of novel anti-infectives. *Microbiome* 8:85. <https://doi.org/10.1186/s40168-020-00866-1>
80. Wieland Brown LC, Acker MG, Clardy J, Walsh CT, Fischbach MA (2009) Thirteen posttranslational modifications convert a 14-residue peptide into the antibiotic thiocillin. *Proc Natl Acad Sci USA* 106:2549–2553. <https://doi.org/10.1073/pnas.0900008106>
81. Rhoads A, Au KF (2015) PacBio sequencing and its applications. *Genom Proteom Bioinform* 13:278–289. <https://doi.org/10.1016/j.gpb.2015.08.002>
82. Lu H, Giordano F, Ning Z (2016) Oxford nanopore MinION sequencing and genome assembly. *Genom Proteom Bioinform* 14:265–279. <https://doi.org/10.1016/j.gpb.2016.05.004>
83. Lozano C, Rezusta A, Ferrer I, Pérez-Laguna V, Zarazaga M, Ruiz-Ripa L, Revillo MJ, Torres C (2017) *Staphylococcus pseud-intermedius* human infection cases in Spain: dog-to-human transmission. *Vector-Borne Zoonotic Dis* 17(4):268–270. <https://doi.org/10.1089/vbz.2016.2048>
84. Gómez P, Casado C, Sáenz Y, Ruiz-Ripa L, Estepa V, Zarazaga M, Torres C (2017) Diversity of species and antimicrobial resistance determinants of staphylococci in superficial waters in Spain. Smalla K, editor. *FEMS Microbiol Ecol* 93(1):fiw208. <https://doi.org/10.1093/femsec/fiw208>
85. Ruiz-Ripa L, Alcalá L, Simón C, Gómez P, Mama OM, Rezusta A, Zarazaga M, Torres C (2019) Diversity of *Staphylococcus aureus* clones in wild mammals in Aragon, Spain, with detection of MRSA ST130-*mecC* in wild rabbits. *J Appl Microbiol* 127(1):284–291. <https://doi.org/10.1111/jam.14301>
86. Ruiz-Ripa L, Gómez P, Alonso CA, Camacho MC, Ramiro Y, de la Puente J, Fernández-Fernández R, Quevedo MA, Blanco JM, Báguena G, Zarazaga M, Höfle U, Torres C (2020) Frequency and characterization of antimicrobial resistance and virulence genes of coagulase-negative staphylococci from wild birds in

- Spain. Detection of *tst*-carrying *S. sciuri* isolates. *Microorganisms* 8(9):1317. <https://doi.org/10.3390/microorganisms8091317>
87. Mama OM, Ruiz-Ripa L, Lozano C, González-Barrio D, Ruiz-Fons JF, Torres C (2019) High diversity of coagulase negative staphylococci species in wild boars, with low antimicrobial resistance rates but detection of relevant resistance genes. *Comp Immunol*

Microbiol Infect Dis 64:125–129. <https://doi.org/10.1016/j.cimid.2019.03.006>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.